



PHE publications gateway number: GW-774

Low-dose diphtheria, tetanus and inactivated poliomyelitis vaccine (Td/IPV) Patient Group Direction (PGD)

This PGD is for the administration of low-dose diphtheria, tetanus and inactivated poliomyelitis vaccine (Td/IPV) to individuals from 10 years of age, in accordance with the national immunisation programme, for travel, or for the management of cases and contacts of diphtheria, tetanus or poliomyelitis.

This PGD is for the administration of Td/IPV by registered healthcare practitioners identified in Section 3, subject to any limitations to authorisation detailed in Section 2.

Reference no: Td/IPV (Revaxis®) PGD
Version no: v03.00
Valid from: 01 November 2019
Review date: 1 May 2021
Expiry date: 31 October 2021

Public Health England has developed this PGD to facilitate publicly-funded immunisation in line with national recommendations.

Those using this PGD must ensure that it is organisationally authorised and signed in Section 2 by an appropriate authorising person, relating to the class of person by whom the product is to be supplied, in accordance with Human Medicines Regulations 2012 (HMR2012)¹. **The PGD is not legal or valid without signed authorisation in accordance with [HMR2012 Schedule 16 Part 2](#).**

Authorising organisations must not alter, amend or add to the clinical content of this document (sections 4, 5 and 6); such action will invalidate the clinical sign-off with which it is provided. In addition, authorising organisations must not alter section 3 'Characteristics of staff'. Only sections 2 and 7 can be amended within the designated editable fields provided.

Operation of this PGD is the responsibility of commissioners and service providers. The final authorised copy of this PGD should be kept by the authorising organisation completing Section 2 for 8 years after the PGD expires if the PGD relates to adults only and for 25 years after the PGD expires if the PGD relates to children only, or adults and children. Provider organisations adopting authorised versions of this PGD should also retain copies for the periods specified above.

Individual practitioners must be authorised by name, under the current version of this PGD before working according to it.

Practitioners and organisations must check that they are using the current version of the PGD. Amendments may become necessary prior to the published expiry date. Current versions of PHE PGD templates for authorisation can be found from:

<https://www.gov.uk/government/collections/immunisation-patient-group-direction-pgd>

Any concerns regarding the content of this PGD should be addressed to:
immunisation@phe.gov.uk




¹ This includes any relevant amendments to legislation (eg [2013 No.235](#), [2015 No.178](#) and [2015 No.323](#)).
Td/IPV (Revaxis®) PGD v03.00 Valid from: 01/11/2019 Expiry: 31/10/2021

Change history

Version number	Change details	Date
V01.00	New PHE PGD template	16 October 2015
V02.00	Td/IPV (Revaxis [®]) PGD routine review and amended to: <ul style="list-style-type: none"> • include vaccination in line with recommendations for the management of diphtheria or polio • remove exclusions regarding timing of previous vaccination (see dose section for schedules) • remove exclusions relating to neurological conditions, encephalopathy and Guillain Barre/brachial neuritis and relevant advice moved to the cautions section • update off-label section in relation to amended indications • update dose section with management of cases and contacts of polio and diphtheria • include minor rewording, layout and formatting changes for clarity and consistency with other PHE PGD templates 	29 September 2017
V03.00	Td/IPV (Revaxis [®]) PGD routine review and amended to: <ul style="list-style-type: none"> • include minor rewording, layout and formatting changes for clarity and consistency with other PHE PGD templates 	12 September 2019

1. PGD development

This PGD has been developed by the following health professionals on behalf of Public Health England:

Developed by:	Name	Signature	Date
Pharmacist (Lead Author)	Elizabeth Graham Lead Pharmacist Immunisation Services, PHE		16/09/2019
Doctor	Mary Ramsay Consultant Epidemiologist and Head of Immunisation and Countermeasures, PHE		16/09/2019
Registered Nurse (Chair of Expert Panel)	David Green Nurse Consultant, Immunisation and Countermeasures, PHE		16/09/2019

This PGD has been peer reviewed by the PHE Immunisations PGD Expert Panel in accordance with PHE PGD Policy. It has been ratified by the PHE Medicines Management Group and the PHE Quality and Clinical Governance Delivery Board.

Expert Panel

Name	Designation
Gayatri Amirthalingam	Consultant Epidemiologist, Public Health England
Ed Gardner	Advanced Paramedic Practitioner/Emergency Care Practitioner, Medicines Manager, Proactive Care Lead
Michelle Jones	Senior Medicines Optimisation Pharmacist, NHS Bristol North Somerset & South Gloucestershire CCG
Jacqueline Lamberty	Lead Pharmacist Medicines Management Services, Public Health England
Vanessa MacGregor	Consultant in Communicable Disease Control, Public Health England, East Midlands Health Protection Team
Alison Mackenzie	Consultant in Public Health Medicine, Screening and Immunisation Lead, Public Health England (South West) / NHS England and NHS Improvement South (South West)
Gill Marsh	Senior Screening and Immunisation Manager, Public Health England / NHS England and NHS Improvement Lancashire and South Cumbria
Lesley McFarlane	Screening and Immunisation Co-ordinator, Public Health England / NHS England and NHS Improvement Leicestershire, Lincolnshire and Northamptonshire
Tushar Shah	Pharmacy Advisor, NHS England and NHS Improvement London Region
Sharon Webb	Programme Manager / Registered Midwife, NHS Infectious Diseases in Pregnancy Screening Programme, Public Health England


2. Organisational authorisations




The PGD is not legally valid until it has had the relevant organisational authorisation.

It is the responsibility of the organisation that has legal authority to authorise the PGD, to ensure that all legal and governance requirements are met. The authorising body accepts governance responsibility for the appropriate use of the PGD.

NHS England and NHS Improvement East of England authorises this PGD for use by the services or providers listed below:

Authorised for use by the following organisations and/or services
All NHS England and NHS Improvement East of England commissioned immunisation services or NHS Trust providing immunisation services covering Norfolk, Suffolk, Cambridgeshire, Peterborough, Essex, Southend-on-Sea, Thurrock, Bedfordshire, Hertfordshire, Luton and Milton Keynes local authorities, and Health and Justice facilities where NHS England and NHS Improvement East of England is the commissioner.
Limitations to authorisation
None

Organisational approval (legal requirement)			
Role	Name	Sign	Date
Deputy Medical Director	Dr. James Hickling		10/10/2019

Additional signatories according to locally agreed policy			
Role	Name	Sign	Date
Screening and Immunisation Lead	Dr. Pam Hall		09/10/2019
Pharmacist	Dr. Paul Duell		07/10/2019
Screening and Immunisation Coordinator	Alex Burghlea		01/10/2019

Local enquiries regarding the use of this PGD may be directed to

For East Anglia email: England.ea-phsi@nhs.net

For Essex email: England.essexatimms@nhs.net

For Bedfordshire, Hertfordshire, Luton and Milton Keynes email: England.immsga@nhs.net

Section 7 provides a practitioner authorisation sheet. Individual practitioners must be authorised by name to work to this PGD. Alternative practitioner authorisation sheets may be used where appropriate in accordance with local policy but this should be an individual agreement or a multiple practitioner authorisation sheet as included at the end of this PGD.

3. Characteristics of staff

<p>Qualifications and professional registration</p>	<p>Registered professional with one of the following bodies:</p> <ul style="list-style-type: none"> nurses and midwives currently registered with the Nursing and Midwifery Council (NMC) pharmacists currently registered with the General Pharmaceutical Council (GPhC) (Note: This PGD is not relevant to privately provided community pharmacy services) paramedics and physiotherapists currently registered with the Health and Care Professions Council (HCPC) <p>The practitioners above must also fulfil the Additional requirements detailed below.</p> <p>Check Section 2 Limitations to authorisation to confirm whether all practitioners listed above have organisational authorisation to work under this PGD.</p>
<p>Additional requirements</p>	<p>Additionally, practitioners:</p> <ul style="list-style-type: none"> must be authorised by name as an approved practitioner under the current terms of this PGD before working to it must have undertaken appropriate training for working under PGDs for supply/administration of medicines must be competent in the use of PGDs (see NICE Competency framework for health professionals using PGDs) must be familiar with the vaccine product and alert to changes in the Summary of Product Characteristics (SPC), Immunisation Against Infectious Disease (‘The Green Book’), and national and local immunisation programmes must have undertaken training appropriate to this PGD as required by local policy and in line with the National Minimum Standards and Core Curriculum for Immunisation Training must be competent to undertake immunisation and to discuss issues related to immunisation must be competent in the handling and storage of vaccines, and management of the ‘cold chain’ must be competent in the recognition and management of anaphylaxis must have access to the PGD and associated online resources should fulfil any additional requirements defined by local policy <p>The individual practitioner must be authorised by name, under the current version of this PGD before working according to it.</p>
<p>Continued training requirements</p>	<p>Practitioners must ensure they are up to date with relevant issues and clinical skills relating to immunisation and management of anaphylaxis, with evidence of appropriate Continued Professional Development (CPD).</p> <p>Practitioners should be constantly alert to any subsequent recommendations from Public Health England and/or NHS England and other sources of medicines information.</p> <p>Note: The most current national recommendations should be followed but a Patient Specific Direction (PSD) may be required to administer the vaccine in line with updated recommendations that are outside the criteria specified in this PGD.</p>

4. Clinical condition or situation to which this PGD applies

<p>Clinical condition or situation to which this PGD applies</p>	<p>Indicated for the active immunisation of individuals from 10 years of age for the prevention of diphtheria, tetanus and poliomyelitis, in accordance with the national immunisation programme and recommendations given in Chapter 15, Chapter 26 and Chapter 30 of Immunisation Against Infectious Disease: 'The Green Book'.</p>
<p>Criteria for inclusion</p>	<p>Individuals aged 10 years and over who:</p> <ul style="list-style-type: none"> • require a booster following a primary course of immunisation against diphtheria, tetanus and poliomyelitis (this booster is usually offered at 13 to 18 years of age, unless the course has already been completed) • have no history or an incomplete history of diphtheria, tetanus or poliomyelitis immunisation • are travelling to an area where medical attention may not be accessible should a tetanus prone wound occur, or will be residing in epidemic or endemic areas where tetanus, diphtheria or poliomyelitis protection is required, and the final dose of the relevant antigen was received more than 10 years ago, even if the individual has received 5 doses of tetanus containing vaccine previously • have a tetanus prone wound and one or more of the following apply (see Green Book Chapter 30): <ul style="list-style-type: none"> ○ primary tetanus immunisation is incomplete ○ tetanus boosters are not up to date or last dose of tetanus containing vaccine was more than 10 years ago ○ tetanus immunisation status is unknown or uncertain ○ individual has never received tetanus immunisation • require vaccination in line with recommendations for the management of cases and contacts of diphtheria or polio
<p>Criteria for exclusion²</p>	<p>Individuals for whom no valid consent has been received.</p> <p>Individuals who:</p> <ul style="list-style-type: none"> • are aged less than 10 years • have had a confirmed anaphylactic reaction to a previous dose of diphtheria, tetanus or poliomyelitis containing vaccine, including any conjugate vaccines where diphtheria or tetanus toxoid is used in the conjugate • have had a confirmed anaphylactic reaction to any component of the vaccine, including neomycin, streptomycin or polymyxin B • are suffering from acute severe febrile illness (the presence of a minor infection is not a contraindication for immunisation)
<p>Cautions including any relevant action to be taken</p> <p>Continued over page</p>	<p>Td/IPV may be given to pregnant women when protection is required without delay, such as following a tetanus prone wound. However, pregnant women from week 16 of pregnancy onwards should instead be protected by the administration of the routinely indicated dTaP/IPV (see Pertussis PGD).</p> <p>The presence of a neurological condition is not a contraindication to immunisation but if there is evidence of current neurological</p>

² Exclusion under this PGD does not necessarily mean the medication is contraindicated, but it would be outside its remit and another form of authorisation will be required

<p>Cautions including any relevant action to be taken (continued)</p>	<p>deterioration, deferral of vaccination may be considered, to avoid incorrect attribution of any change in the underlying condition. The risk of such deferral should be balanced against the risk of the preventable infection, and vaccination should be promptly given once the diagnosis and/or the expected course of the condition becomes clear.</p> <p>If a child has experienced encephalopathy or encephalitis within seven days of immunisation, it is unlikely that these conditions will have been caused by the vaccine and they should be investigated by a specialist. If a cause is identified or the child recovered within seven days, immunisation should proceed as recommended. In children where no underlying cause was found, and the child did not recover completely within seven days, immunisation should be deferred until the condition has stabilized or the expected course of the condition becomes clear.</p> <p>The immunogenicity of the vaccine could be reduced in immunosuppressed subjects. Where possible, vaccination should be postponed until immune function has recovered. However, vaccination of subjects with chronic immunodeficiency, such as AIDS, is recommended even if the antibody response might be limited.</p>
<p>Action to be taken if the patient is excluded</p>	<p>If aged under 10 years assess for immunisation with DTaP/IPV/Hib/HepB, DTaP/IPV or dTaP/IPV as appropriate.</p> <p>In case of postponement due to acute severe febrile illness, advise when the individual can be vaccinated and ensure another appointment is arranged.</p> <p>Seek appropriate advice from the local Screening and Immunisation Team, local Health Protection Team or the individual's clinician as appropriate.</p> <p>The risk to the individual of not being immunised must be taken into account.</p> <p>Document the reason for exclusion and any action taken in the individual's clinical records.</p> <p>Inform or refer to the GP or a prescriber as appropriate.</p>
<p>Action to be taken if the patient or carer declines treatment</p>	<p>Informed consent, from the individual or a person legally able to act on the person's behalf, must be obtained for each administration.</p> <p>Advise the individual/parent/carer about the protective effects of the vaccine, the risks of infection and potential complications.</p> <p>Document advice given and the decision reached.</p> <p>Inform or refer to the GP or a prescriber as appropriate.</p>
<p>Arrangements for referral for medical advice</p>	<p>As per local policy</p>

5. Description of treatment

Name, strength & formulation of drug	Adsorbed diphtheria (low dose), tetanus, and inactivated poliomyelitis vaccine (Td/IPV): <ul style="list-style-type: none"> • Revaxis[®], suspension for injection in a pre-filled syringe.
Legal category	Prescription Only Medicine (POM)
Black triangle▼	No
Off-label use	<p>Primary immunisation is off-label administration in accordance with the recommendations given for individuals over 10 years of age in Chapter 15, Chapter 26 and Chapter 30 of Immunisation Against Infectious Disease: 'The Green Book'.</p> <p>Administration to individuals who have received a vaccine containing diphtheria or tetanus toxoids within the previous five years is off-label but indicated for the management of primary immunisation (as above) and for cases and contacts of diphtheria or polio in accordance with PHE disease management guidelines (see Dose and frequency of administration).</p> <p>Administration to individuals who experienced neurological complications following an earlier immunisation against diphtheria and/or tetanus is off-label but may proceed once the cause is identified, the condition has been stabilized or the expected course of the condition becomes clear in accordance with the recommendations in Chapter 15 and Chapter 30 of Immunisation Against Infectious Disease: 'The Green Book'.</p> <p>Vaccine should be stored according to the conditions detailed in the Storage section below. However, in the event of an inadvertent or unavoidable deviation of these conditions refer to PHE Vaccine Incident Guidance. Where vaccine is assessed in accordance with these guidelines as appropriate for continued use this would constitute off-label administration under this PGD.</p> <p>Where a vaccine is recommended off-label consider, as part of the consent process, informing the individual/parent/carer that the vaccine is being offered in accordance with national guidance but that this is outside the product licence.</p>
Route / method of administration	<p>Administer by intramuscular injection, preferably into deltoid region of the upper arm.</p> <p>When administering at the same time as other vaccines care should be taken to ensure that the appropriate route of injection is used for all the vaccinations.</p> <p>The vaccines should be given at separate sites, preferably in different limbs. If given in the same limb, they should be given at least 2.5cm apart. The site at which each vaccine was given should be noted in the individual's records.</p> <p>For individuals with a bleeding disorder, vaccines normally given by an intramuscular route should be given in accordance with the recommendations in 'The Green Book' Chapter 4.</p>

Continued over page

<p>Route / method of administration (continued)</p>	<p>The vaccine's normal appearance is a cloudy white suspension that may sediment during storage. Shake the pre-filled syringe well to distribute uniformly the suspension before administering the vaccine.</p> <p>The vaccine should not be used if foreign particles are present in the suspension.</p> <p>The SPC provides further guidance on administration and is available from the electronic Medicines Compendium website: www.medicines.org.uk</p>
<p>Dose and frequency of administration</p> <p>Continued over page</p>	<p>Single 0.5ml dose per administration</p> <p>Routine childhood immunisation schedule</p> <p>Td/IPV is routinely offered to teenagers as a second booster dose at around 14 years of age. It should ideally be given 10 years after the first booster dose. It should be given at the school session or scheduled appointment provided a minimum of 5 years have elapsed between the first and second boosters. (Note: the first booster is usually given at pre-school age using dTaP/IPV or DTaP/IPV (Repevax[®] or Infanrix[®]-IPV)).</p> <p>UK immunisation schedule for previously unimmunised individuals or where there is an unknown or incomplete history of diphtheria, tetanus and poliomyelitis vaccination</p> <p>Infants with uncertain or incomplete diphtheria, tetanus and poliomyelitis vaccine history should be vaccinated in accordance with the vaccination of individuals with uncertain or incomplete immunisation status flow chart.</p> <p>The primary course consists of three doses, allowing an interval of one month between doses. Where a primary course is interrupted it should be resumed but not repeated.</p> <p>A first booster dose should be administered at least 5 years after the third dose of the primary course.</p> <p>A second booster dose should be administered a minimum of 5 years, ideally 10 years, after the first booster dose, if less than 5 doses of diphtheria, tetanus and polio vaccine are documented.</p> <p>Travel immunisation</p> <p>Individuals travelling should be vaccinated in accordance with the UK schedule.</p> <p>A single booster dose may be indicated for fully immunised individuals whose last dose of vaccine was more than 10 years ago (see https://travelhealthpro.org.uk/).</p> <p>Management of tetanus prone wound</p> <p>Individuals with a tetanus prone wound who received their last dose of tetanus containing vaccine more than 10 years ago should receive a reinforcing dose of vaccine.</p> <p>Individuals with incomplete or uncertain history of tetanus immunisation should be vaccinated in accordance with the recommendations in the 'Green Book' Chapter 30 Table 30.1.</p> <p>Individuals may also require human tetanus immunoglobulin (see 'Green Book' Chapter 30). Administration of tetanus immunoglobulin is not covered by this PGD.</p>

<p>Dose and frequency of administration (continued)</p>	<p>Management of cases and contacts of diphtheria</p> <p>Cases and contacts of diphtheria should be managed in accordance with Public health control and management of diphtheria (in England and Wales) guidelines and recommendations from the local health protection team.</p> <p>Individuals should have their immunisation status checked to ensure they are up to date with the recommended UK immunisation programmes.</p> <p>Unimmunised individuals should receive three doses at monthly intervals.</p> <p>Individuals who are fully immunised but have not received diphtheria containing vaccine in last 12 months may be given a single reinforcing dose of Td/IPV.</p> <p>Management of cases and contacts of polio</p> <p>Cases and contacts of polio should be managed in accordance with PHE national polio guidelines: Local and regional services and recommendations from the local health protection team.</p> <p>Individuals should have their immunisation status checked to ensure they are up to date with the recommended UK immunisation programmes.</p> <p>Management will depend on the level of exposure but may include the administration of a single dose of IPV containing vaccine, regardless of vaccine history.</p> <p>Points to note</p> <p>Where there is no reliable history of previous immunisation, it should be assumed that individuals are unimmunised and the full UK recommendations should be followed.</p> <p>Where children have had a fourth dose of tetanus, diphtheria and polio containing vaccine at around 18 months of age, this dose should be discounted as it may not provide satisfactory protection until the time of the teenage booster. The routine pre-school and subsequent boosters should be given according to the UK schedule.</p> <p>If a person attends for a routine booster dose and has a history of receiving a vaccine following a tetanus-prone wound, attempts should be made to identify which vaccine was given. If the vaccine given at the time of the injury was the same as that due at the current visit and was given after an appropriate interval, then the routine booster dose is not required. Otherwise, the dose given at the time of injury should be discounted as it may not provide long-term protection against all antigens, and the scheduled immunisation should be given. Such additional doses are unlikely to produce an unacceptable rate of reactions.</p>
<p>Duration of treatment</p> <p>Continued over page</p>	<p>A total of 5 doses (3 primary course doses and 2 booster doses) of diphtheria, tetanus and polio vaccine are indicated for complete immunisation (doses provided under the age of 10 years will not be provided using this vaccine).</p> <p>A further booster dose may be indicated 10 years after the final dose where risk of exposure is high.</p>

Duration of treatment continued	A reinforcing dose may be recommended following potential exposure.
Quantity to be supplied / administered	Single 0.5ml dose per administration.
Supplies	<p>Centrally purchased vaccines for the national immunisation programme for the NHS can only be ordered via ImmForm. Vaccines for use for the national immunisation programme are provided free of charge.</p> <p>Vaccine for indications other than the national immunisation programme should be obtained from manufacturers/wholesalers.</p> <p>Protocols for the ordering, storage and handling of vaccines should be followed to prevent vaccine wastage (see Green Book Chapter 3).</p>
Storage	<p>Store at +2°C to +8°C. Store in original packaging to protect from light. Do not freeze.</p> <p>In the event of an inadvertent or unavoidable deviation of these conditions vaccine that has been stored outside the conditions stated above should be quarantined and risk assessed for suitability of continued off-label use or appropriate disposal, refer to PHE Vaccine Incident Guidance.</p>
Disposal	Equipment used for immunisation, including used vials, ampoules, or discharged vaccines in a syringe or applicator, should be disposed of safely in a UN-approved puncture-resistant 'sharps' box, according to local authority regulations and guidance in the technical memorandum 07-01 : Safe management of healthcare waste (Department of Health, 2013).
Drug interactions	<p>Immunological response may be diminished in those receiving immunosuppressive treatment. Vaccination is recommended even if the antibody response may be limited.</p> <p>May be given at the same time as other vaccines.</p> <p>A detailed list of drug interactions is available in the SPC, which is available from the electronic Medicines Compendium website: www.medicines.org.uk</p>
Identification & management of adverse reactions	<p>Local reactions following vaccination are very common such as pain, swelling or redness at the injection site. A small painless nodule may form at the injection site.</p> <p>Common adverse reactions include pyrexia, headache, vertigo, nausea and vomiting.</p> <p>Allergic reactions can occur including generalised skin reactions such as urticaria, anaphylactic reactions, angioedema and shock.</p> <p>A detailed list of adverse reactions is available in the SPC, which is available from the electronic Medicines Compendium website: www.medicines.org.uk</p>
Reporting procedure of adverse reactions Continued over page	Healthcare professionals and patients/carers are encouraged to report suspected adverse reactions to the Medicines and Healthcare products Regulatory Agency (MHRA) using the Yellow

Reporting procedure of adverse reactions continued	Card reporting scheme (http://yellowcard.mhra.gov.uk). Any adverse reaction to a vaccine should be documented in the individual's record and the individual's GP should be informed.
Written information to be given to patient or carer	Offer marketing authorisation holder's patient information leaflet (PIL) provided with the vaccine.
Patient advice / follow up treatment	Inform individual/carer/parent of possible side effects and their management. The individual/carer/parent should be advised to seek medical advice in the event of an adverse reaction. When administration is postponed advise the individual/carer/parent when to return for vaccination.
Special considerations / additional information	Ensure there is immediate access to adrenaline (epinephrine) 1 in 1000 injection and access to a telephone at the time of vaccination. Minor illnesses without fever or systemic upset are not valid reasons to postpone immunisation. If an individual is acutely unwell, immunisation may be postponed until they have fully recovered. Intravenous drug users are at greater risk of tetanus. Every opportunity should be taken to ensure that they are fully protected against tetanus. Booster doses should be given if there is any doubt about their immunisation status.
Records	Record: <ul style="list-style-type: none"> • that valid informed consent was given • name of individual, address, date of birth and GP with whom the individual is registered • name of immuniser • name and brand of vaccine • date of administration • dose, form and route of administration of vaccine • quantity administered • batch number and expiry date • anatomical site of vaccination • advice given, including advice given if excluded or declines immunisation • details of any adverse drug reactions and actions taken • supplied via PGD Records should be signed and dated (or a password-controlled immuniser's record on e-records). All records should be clear, legible and contemporaneous. This information should be recorded in the individual's GP record. Where vaccine is administered outside the GP setting appropriate health records should be kept and the individual's GP informed. The local Child Health Information Systems team (Child Health Records Department) must be notified using the appropriate documentation/pathway as required by any local or contractual arrangement. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes in accordance with local policy.

6. Key references

Key references	<p>Td/IPV vaccine (Revaxis®)</p> <ul style="list-style-type: none">• Immunisation against infectious disease: The Green Book Chapter 15, Chapter 26 updated 19 April 2013 and Chapter 30 updated 26 November 2018. https://www.gov.uk/government/collections/immunisation-against-infectious-disease-the-green-book• Summary of product characteristic for Revaxis®, Sanofi Pasteur. 4 January 2018. www.medicines.org.uk http://www.medicines.org.uk/emc/medicine/15259• NHS public health functions agreement 2018-19, Service Specification No.12. Td/IPV (teenage booster) immunisation programme. https://www.england.nhs.uk/commissioning/pub-hlth-res/• Vaccination of individuals with uncertain or incomplete immunisation status. Public Health England. Updated 22 August 2019. https://www.gov.uk/government/publications/vaccination-of-individuals-with-uncertain-or-incomplete-immunisation-status• Public health control and management of diphtheria (in England and Wales) guidelines. Public Health England. 24 March 2015. https://www.gov.uk/government/publications/diphtheria-public-health-control-and-management-in-england-and-wales• PHE national polio guidelines: Local and regional services. Public Health England. 4 February 2016. https://www.gov.uk/government/publications/polio-national-guidelines <p>General</p> <ul style="list-style-type: none">• Health Technical Memorandum 07-01: Safe Management of Healthcare Waste. Department of Health 20 March 2013. https://www.gov.uk/government/publications/guidance-on-the-safe-management-of-healthcare-waste• National Minimum Standards and Core Curriculum for Immunisation Training. Published February 2018. https://www.gov.uk/government/publications/national-minimum-standards-and-core-curriculum-for-immunisation-training-for-registered-healthcare-practitioners• NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017. https://www.nice.org.uk/guidance/mpg2• NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017. https://www.nice.org.uk/guidance/mpg2/resources• PHE Immunisation Collection https://www.gov.uk/government/collections/immunisation• PHE Vaccine Incident Guidance https://www.gov.uk/government/publications/vaccine-incident-guidance-responding-to-vaccine-errors
-----------------------	--

7. Practitioner authorisation sheet

Td/IPV (Revaxis®) PGD v03.00 Valid from: 01/11/2019 Expiry: 31/10/2021

Before signing this PGD, check that the document has had the necessary authorisations in section two. Without these, this PGD is not lawfully valid.

Practitioner

By signing this PGD you are indicating that you agree to its contents and that you will work within it.

PGDs do not remove inherent professional obligations or accountability.

It is the responsibility of each professional to practise only within the bounds of their own competence and professional code of conduct.

I confirm that I have read and understood the content of this PGD and that I am willing and competent to work to it within my professional code of conduct.			
Name	Designation	Signature	Date

Authorising manager

I confirm that the practitioners named above have declared themselves suitably trained and competent to work under this PGD. I give authorisation on behalf of INSERT NAME OF ORGANISATION: for the above-named healthcare professionals who have signed the PGD to work under it.			
Name	Designation	Signature	Date

Note to authorising manager

Score through unused rows in the list of practitioners to prevent practitioner additions post managerial authorisation.

This authorisation sheet should be retained to serve as a record of those practitioners authorised to work under this PGD.