
CVD PREMATURE MORTALITY

CVD PREMATURE MORTALITY

CCG Name	2014
LEICESTER CITY CCG	90.1
NOTTINGHAM CITY CCG	89.8
CORBY CCG	89.1
LINCOLNSHIRE WEST CCG	73.5
LINCOLNSHIRE EAST CCG	71.0
MANSFIELD AND ASHFIELD CCG	68.7
HARDWICK CCG	67.2
NEWARK AND SHERWOOD CCG	66.5
ENGLAND	63.7
NORTH DERBYSHIRE CCG	62.8
SOUTHERN DERBYSHIRE CCG	62.8
EREWASH CCG	62.2
NOTTINGHAM NORTH AND EAST CCG	61.3
SOUTH LINCOLNSHIRE CCG	60.8
NENE CCG	58.7
WEST LEICESTERSHIRE CCG	57.0
EAST LEICESTERSHIRE AND RUTLAND CCG	53.8
NOTTINGHAM WEST CCG	53.7
SOUTH WEST LINCOLNSHIRE CCG	49.0
RUSHCLIFFE CCG	35.7

CVD PREMATURE MORTALITY RATE 2009-2014




NUMBER OF PREMATURE DEATHS DUE TO CVD IN 2014

LEADING CAUSES OF YEARS OF LIFE LOST IN ENGLAND **GLOBAL BURDEN OF DISEASE STUDY 2013**

_		
1	Ischaemic heart disease	
2	Lung cancer	
3	Cerebrovascular disease	
4	COPD	
5 Alzheimer's disease		
11	Other cardiovascular	

ATRIAL FIBRILLATION

STROKE ADMISSIONS WITH **KNOWN AF 2014-15**



NUMBER OF

UNDIAGNOSED AF PATIENTS...

% observed to expected **AF** prevalence

High risk AF patients **NOT** on anticoagulation (17.2%)

*includes exception reported

CCG EXCEPTION REPORTING FOR **ANTICOAGULATION**

PRACTICE RANGE OF **EXCEPTION REPORTING**



BLOOD PRESSURE

NUMBER OF PEOPLE WITH **UNDIAGNOSED HYPERTENSION**



observed to expected **HYPERTENSION PREVALENCE**

Actual = 12.0%

Estimated = 23.7%

1.425

Hypertensive patients with BP **NOT TO** QOF target 150/90 (16.1%)

*includes exception reported

HEART ATTACKS IN PEOPLE

UNDER 50 - 2012-13 to 2014/15

CHRONIC KIDNEY DISEASE

NUMBER OF PEOPLE WITH **UNDIAGNOSED CKD**



observed to expected **CKD PREVALENCE**

Actual = 4.0%

Estimated = 5.5%

437

CKD patients with BP **NOT TO** QOF target 140/85 (19.3%)

*includes exception reported

Estimated people with FH

Rate = 1 per 500 Rate = 1 per 250

Estimated 15% diagnosed with FH

Rate = 1 per 500 Rate = 1 per 250

33.1

31.5

28.9

28.9

29.6

32.0

Compared to England average

Smoking prevalence in adults in

current smokers (2014)

routine and manual occupations -

FAMILIAL

HYPERCHOLESTEROLEMIA (FH)

SMOKING DATA BY LOCAL AUTHORITY

19.9

20.7

24.2

18.8

Smoking Prevalence in adults

Derby

Derbyshire

Leicester

Lincolnshire

Nottingham

Rutland

Leicestershire

Northamptonshire

Nottinghamshire

East Midlands

current smokers (2014)

HEART FAILURE PATIENTS

HEART

FAILURE

March 03 2014 March 2015

WITH LVSD

HEART FAILURE ADMISSIONS

90
434
014-15

ſ	Admission Rate per	2014-15	F Mids Av
	Secondary Diagnosis	£1,274,000	£1,242,000
	Primary Diagnosis#	£355,000	£302,000

Admission Rate per 1000 HF Patients	2014-15	E.Mids Av
Primary Diagnosis#	154	179
Secondary Diagnosis	741	749

......

42 Heart Failure LVSD patients **NOT** on dual therapy ACE inhibitor/ARB and a betablocker (29.4%)

DIABETIC **FOOT AMPUTATIONS**



Minor

2,727

11

DIABETES



Practice Participation in 80% **National Diabetes Audit**

2014-15

CARE (O)**PROCESSES** 0

TYPE 1 - 54.0% England = 39.6% E.Mids CCG Range = 29.3 - 54.0%


Lowest 8 Care Process 70.9% Urine Albumin

TYPE 2 - 62.3% England = 59.0% E.Mids CCG Range = 51.2 - 73.8%

Lowest 8 Care Process

% OF DIABETIC PATIENTS **MEETING 3 TARGETS** 2014-15

TYPE 1 - 19.8% England = 19.3% E.Mids CCG Range :



Produced by: **East Midlands Clinical Networks**

E.Mids CCG Range = 35.6 - 44.6%

East Midlands Rate in 2011 - 2014 Total - 2.37 Major - 0.82 Minor - 1.55

1.14