

NHS Lincolnshire East CCG

Diabetic Foot Amputations

Number of Amputations

	Total Amputations	Major Amputations	Minor Amputations																		
2009 - 12	124	49	75																		
2010 - 13	133	50	83																		
2011 - 14	144	51	93																		
2012 - 15	135	41	94																		
	<table><tr><td>2012-13</td><td>44</td></tr><tr><td>2013-14</td><td>48</td></tr><tr><td>2014-15</td><td>43</td></tr></table>	2012-13	44	2013-14	48	2014-15	43	<table><tr><td>2012-13</td><td>15</td></tr><tr><td>2013-14</td><td>15</td></tr><tr><td>2014-15</td><td>11</td></tr></table>	2012-13	15	2013-14	15	2014-15	11	<table><tr><td>2012-13</td><td>29</td></tr><tr><td>2013-14</td><td>33</td></tr><tr><td>2014-15</td><td>32</td></tr></table>	2012-13	29	2013-14	33	2014-15	32
2012-13	44																				
2013-14	48																				
2014-15	43																				
2012-13	15																				
2013-14	15																				
2014-15	11																				
2012-13	29																				
2013-14	33																				
2014-15	32																				

Amputation Rate per 1,000 adults with diabetes

	Total Amputations	Major Amputations	Minor Amputations
2009 - 12	2.8 2.6 England	1.1 1.0 England	1.7 1.1 England
2010 - 13	2.9 2.6 England	1.1 0.9 England	1.8 1.7 England
2011 - 14	3.0 2.6 England	1.1 0.8 England	1.9 1.8 England
2012 - 15	2.7 2.6 England	0.8 0.8 England	1.9 1.8 England

Hospital Stays for Diabetic Foot Disease



	Episodes of care			Nights in hospital			Average LOS	
	Number	Rate per 1,000		Number	Rate per 1,000			
2009 - 12	817	18.4	18.3 England	8,287	186.9	166.1 England	10.1	9.1 England
2010 - 13	916	19.8	18.6 England	9,579	207.0	165.1 England	10.5	8.9 England
2011 - 14	966	20.0	19.2 England	9,113	188.6	163.2 England	9.4	8.5 England
2012 - 15	932	18.6	19.8 England	7,698	153.4	161.0 England	8.3	8.1 England

Key

- Significantly lower than the England value
- No significant difference from the England value
- Significantly higher than the England value

Produced October 2016 by:



East Midlands Clinical Networks

Data Source: Public Health England

Acknowledgement: Thanks to Public Health England for providing data analysis support