


Shingles vaccine







The shingles vaccine helps protect you from shingles.

Shingles can make you really **poorly** sometimes for many weeks.

You can get shingles more than once.

If you have shingles you might feel poorly first and then a **rash** will appear later.

The shingles rash is **liquid filled** spots which are **red**, **painful** and **can itch or sting**.

The red rash can be harder to see on black or brown skin.

The rash can be on any part of your body but will only be on **one side**.


Shingles may cause you pain even after the rash has

If you have shingles

You may not feel hungry

You might have a painful headache and feel unwell.

You may find it hard to sleep.







You **cannot catch** shingles. You can only get shingles if you have had chickenpox.

If you have shingles, you **can give** chickenpox to someone who has not had chickenpox before.

Not everyone who has had chickenpox will get shingles.

Who can have the shingles vaccine

- Anyone aged 65 on or after 1st September 2023
- Anyone aged **70 79** who has not had the vaccine
- Anyone aged **50 or over** who has a **severely** weakened immune system.

You can have the vaccine even if you have had shingles before.

The shingles vaccine is given as two injections.

The **second** injection is given between **two and twelve** months after the first injection.



Side effects from the injection are usually not serious.

Sometimes you can get a **lump** or **pain** where the injection was given, a **headache**, a **fever**, feeling **tired**, feeling **sick** or being sick or feeling **shivery**.

Not everyone gets side effects.

The vaccine is free safe

It is important to be protected from serious disease.

Phone your GP Practice to book an appointment for your vaccine.





