



Meningococcal Group B Vaccine Patient Group Direction (PGD)

This PGD is for the administration of meningococcal group B vaccine (rDNA, component, adsorbed) (4CMenB) to individuals from 8 weeks of age eligible for the national routine immunisation programme and to individuals for the prevention of secondary cases of meningococcal group B disease.

This PGD is for the administration of 4CMenB by registered healthcare practitioners identified in [Section 3](#), subject to any limitations to authorisation detailed in [Section 2](#).

Reference no: MenB PGD
Version no: v6.00
Valid from: 28 February 2023
Review date: 1 September 2024
Expiry date: 28 February 2025

The UK Health Security Agency (UKHSA) has developed this PGD to facilitate the delivery of publicly funded immunisations in England in line with national recommendations.

Those using this PGD must ensure that it is organisationally authorised and signed in Section 2 by an appropriate authorising person, relating to the class of person by whom the product is to be supplied, in accordance with Human Medicines Regulations 2012 (HMR2012)¹. **The PGD is not legal or valid without signed authorisation in accordance with [HMR2012 Schedule 16 Part 2](#).**

Authorising organisations must not alter, amend or add to the clinical content of this document (sections 4, 5 and 6); such action will invalidate the clinical sign-off with which it is provided. In addition, authorising organisations must not alter section 3 'Characteristics of staff'. Only sections 2 and 7 can be amended within the designated editable fields provided.

Operation of this PGD is the responsibility of commissioners and service providers. The final authorised copy of this PGD should be kept by the authorising organisation completing Section 2 for 8 years after the PGD expires if the PGD relates to adults only and for 25 years after the PGD expires if the PGD relates to children only, or adults and children. Provider organisations adopting authorised versions of this PGD should also retain copies for the periods specified above.

Individual practitioners must be authorised by name, under the current version of this PGD before working according to it.

Practitioners and organisations must check that they are using the current version of the PGD. Amendments may become necessary prior to the published expiry date. Current versions of the UKHSA PGD templates for authorisation can be found from:

<https://www.gov.uk/government/collections/immunisation-patient-group-direction-pgd>

Any concerns regarding the content of this PGD should be addressed to: immunisation@ukhsa.gov.uk

Enquiries relating to the availability of organisationally authorised PGDs and subsequent versions of this PGD should be directed to: your local screening and immunisation team

¹ This includes any relevant amendments to legislation.




Change History

Version number	Change details	Date
V1.00	New MenB PHE PGD Template	21 July 2015
V2.00	<p>PHE MenB PGD amended to:</p> <ul style="list-style-type: none"> • include immunisation into the thigh for individuals over 1 year of age • update dosing recommendations for individuals with incomplete vaccination status • reference the protocol for ordering storage and handling of vaccines • update wording regarding authorisation in line with agreed PHE PGD template changes • include minor rewording, layout and formatting changes for clarity and consistency with other PHE PGD templates 	3 February 2017
V3.00	<p>PHE MenB PGD amended to:</p> <ul style="list-style-type: none"> • update dosing guidance for the prevention of secondary cases of meningococcal group B disease, see Annex A, in line with revised Public Health England Guidance for Public Health Management of Meningococcal Disease in the UK • include additional healthcare practitioners (pharmacists, paramedics, physiotherapists) in Section 3 • refer to the MenB risk groups PGD in the inclusion criteria section • refer to vaccine incident guidelines in off-label and storage sections • include rewording, layout and formatting changes for clarity and consistency with other PHE PGD templates 	24 April 2018
V4.00	<p>PHE MenB PGD amended to:</p> <ul style="list-style-type: none"> • remove the black triangle status • update details regarding permissible use of Immform supplies of 4CMenB • include rewording, layout and formatting changes for clarity and consistency with other PHE PGD templates 	21 December 2018
V5.00	<p>PHE MenB PGD amended to:</p> <ul style="list-style-type: none"> • update off-label section because SPC now includes administration of 2+1 schedule starting at 2 months • update adverse drug reactions section • include a caution relating to immunosuppressed individuals • update adverse drug reactions section • include rewording, layout and formatting changes for clarity and consistency with other PHE PGD templates 	28 January 2021

V6.00	<p>UKHSA MenB PGD amended to:</p> <ul style="list-style-type: none"> • include minor rewording of standard text, layout and formatting changes for clarity and consistency with organisation change and other UKHSA PGDs • amend NHS England and Improvement (NHSE) to NHS England (NHSE) following completion of merger on 1 July 2022 • align the management of anaphylaxis with other UKHSA PGDs in cautions section • add the formulation and strength to the name of the drug • update the advice for individuals with unknown or incomplete history of vaccination in dose and frequency section • include in dose and frequency premature infants, HIV and immunosuppressed cohorts • update drug interactions in accordance with SPC update • update adverse reactions in accordance with updated SPC • update advice for administration of paracetamol in adverse reactions section • update references • remove the table for schedule guidance for secondary prevention of MenB disease as linked in references and through the PGD 	7 December 2022
-------	---	-----------------

1. PGD development

This PGD has been developed by the following health professionals on behalf of the UKHSA:

Developed by:	Name	Signature	Date
Pharmacist (Lead Author)	Suki Hunjunt Lead Pharmacist Immunisation Services, Immunisation and Vaccine Preventable Diseases Division, UKHSA		15 December 2022
Doctor	Mary Ramsay Consultant Epidemiologist, Immunisation and Vaccine Preventable Diseases Division, UKHSA		15 December 2022
Registered Nurse (Chair of Expert Panel)	David Green Nurse Consultant for Immunisation, Immunisation and Vaccine Preventable Diseases Division, UKHSA		15 December 2022

This PGD has been peer reviewed by the UKHSA Immunisations PGD Expert Panel in accordance with the UKHSA PGD Policy. It has been approved by the UKHSA Medicines Governance Group and ratified by the UKHSA Clinical Quality and Oversight Board.

Expert Panel

Nicholas Aigbogun	Consultant in Communicable Disease Control, Yorkshire and Humber Health Protection Team, UKHSA
Gayatri Amrithalingam	Consultant Epidemiologist, Immunisation and Vaccine Preventable Diseases Division, UKHSA
Sarah Dermont	Clinical Project Coordinator and Registered Midwife, NHS Infectious Diseases in Pregnancy Screening Programme, NHS England (NHSE)
Ed Gardner	Advanced Paramedic Practitioner/Emergency Care Practitioner, Medicines Manager, Proactive Care Lead
Jacqueline Lamberty	Lead Pharmacist, Medicines Governance, UKHSA
Michelle Jones	Principal Medicines Optimisation Pharmacist, Bristol North Somerset and South Gloucestershire Integrated Care Board
Shamez Ladhani	Paediatric Infectious Disease Consultant, UKHSA
Elizabeth Lockett	Senior Screening & Immunisation Manager NHSE South West
Vanessa MacGregor	Consultant in Communicable Disease Control, East Midlands Health Protection Team, UKHSA
Alison Mackenzie	Consultant in Public Health Medicine, Screening and Immunisation Lead, NHSE South West
Lesley McFarlane	Lead Immunisation Nurse Specialist Immunisation and Vaccine Preventable Diseases Division, UKHSA
Gill Marsh	Principal Screening and Immunisation Manager, NHSE North West
Tushar Shah	Lead Pharmacy Advisor, NHSE London


2. Organisational authorisations

The PGD is not legally valid until it has had the relevant organisational authorisation.

It is the responsibility of the organisation that has legal authority to authorise the PGD, to ensure that all legal and governance requirements are met. The authorising body accepts governance responsibility for the appropriate use of the PGD.

NHS England South East authorises this PGD for use by the services or providers listed below:

Authorised for use by the following organisations and/or services
All NHS England commissioned immunisation services within the NHS England South East Region.
Limitations to authorisation
This patient group direction (PGD) must only be used by the registered healthcare practitioners identified in Section 3 who have been named by their organisation to practice under it. The most recent in-date final version authorised by NHS England (South East) must be used.
This PGD includes vaccination of individuals across the national immunisation programme. Users of this PGD should note that where they are commissioned to immunise certain groups this PGD does not constitute permission to offer immunisation beyond the groups they are commissioned to immunise.

Organisational approval (legal requirement)			
Role	Name	Sign	Date
South East Medical Director System Improvement and Professional Standards	Dr Alison Taylor		24.02.23

Additional signatories according to locally agreed policy			
Role	Name	Sign	Date

Local enquiries regarding the use of this PGD may be directed to your local screening and immunisation team

Section 7 provides a practitioner authorisation sheet. Individual practitioners must be authorised by name to work to this PGD. Alternative practitioner authorisation sheets may be used where appropriate in accordance with local policy but this should be an individual agreement or a multiple practitioner authorisation sheet as included at the end of this PGD.

3. Characteristics of staff

<p>Qualifications and professional registration</p>	<p>Registered professional with one of the following bodies:</p> <ul style="list-style-type: none"> nurses and midwives currently registered with the Nursing and Midwifery Council (NMC) pharmacists currently registered with the General Pharmaceutical Council (GPhC) (Note: This PGD is not relevant to privately provided community pharmacy services) paramedics and physiotherapists currently registered with Health and Care Professions Council (HCPC) <p>The practitioners above must also fulfil the Additional requirements detailed below.</p> <p>Check Section 2 Limitations to authorisation to confirm whether all practitioners listed above have organisational authorisation to work under this PGD.</p>
<p>Additional requirements</p>	<p>Additionally practitioners:</p> <ul style="list-style-type: none"> must be authorised by name as an approved practitioner under the current terms of this PGD before working to it must have undertaken appropriate training for working under PGDs for supply/administration of medicines must be competent in the use of PGDs (see NICE Competency framework for health professionals using PGDs) must be familiar with the vaccine product and alert to changes in the Summary of Product Characteristics (SPC), Immunisation Against Infectious Disease (‘The Green Book’), and national and local immunisation programmes must have undertaken training appropriate to this PGD as required by local policy and in line with the National Minimum Standards and Core Curriculum for Immunisation Training must be competent to undertake immunisation and to discuss issues related to immunisation must be competent in the handling and storage of vaccines, and management of the cold chain must be competent in the recognition and management of anaphylaxis must have access to the PGD and associated online resources should fulfil any additional requirements defined by local policy <p>The individual practitioner must be authorised by name, under the current version of this PGD before working according to it.</p>
<p>Continued training requirements</p>	<p>Practitioners must ensure they are up to date with relevant issues and clinical skills relating to immunisation and management of anaphylaxis, with evidence of appropriate Continued Professional Development (CPD).</p> <p>Practitioners should be constantly alert to any subsequent recommendations from the UKHSA and/or NHSE and other sources of medicines information.</p> <p>Note: The most current national recommendations should be followed but a Patient Specific Direction (PSD) may be required to administer the vaccine in line with updated recommendations that are outside the criteria specified in this PGD.</p>

4. Clinical condition or situation to which this PGD applies

<p>Clinical condition or situation to which this PGD applies</p>	<p>Indicated for the active immunisation of individuals from 8 weeks of age against <i>Neisseria meningitidis</i> group B and for the prevention of secondary cases of meningococcal group B disease, in accordance with the recommendations given in Chapter 22 of Immunisation Against Infectious Disease: The Green Book and Guidance for Public Health Management of Meningococcal Disease in the UK.</p>
<p>Criteria for inclusion</p>	<p>Individuals who:</p> <ul style="list-style-type: none"> • are aged from 8 weeks up to their second birthday and require routine immunisation • require vaccination for the prevention of secondary cases of Men B, following specific advice from UKHSA Health Protection Teams and in accordance with Guidance for Public Health Management of Meningococcal Disease in the UK. <p>Note: Individuals, from 2 years of age, with an underlying medical condition which puts them at increased risk from <i>Neisseria meningitidis</i> group B, that is individuals with asplenia, splenic dysfunction or complement disorders (including those on, or due to receive, complement inhibitor treatment such as eculizumab), may require additional 'routine' vaccination outside the inclusion criteria for this PGD - see Meningococcal Group B Vaccine Risk Groups PGD and Chapter 7 of 'The Green Book'.</p>
<p>Criteria for exclusion²</p>	<p>Individuals for whom no valid consent has been received.</p> <p>Individuals who:</p> <ul style="list-style-type: none"> • are less than 8 weeks old • are from 2 years of age, unless advised by the UKHSA for the prevention of secondary cases of MenB infection • have had a confirmed anaphylactic reaction to a previous dose of the vaccine • have had a confirmed anaphylactic reaction to any constituent or excipient of the vaccine including kanamycin • require vaccination for occupational health reasons, travel or going to reside abroad • have a history of anaphylactic allergy to latex • are suffering from acute severe febrile illness (the presence of a minor infection is not a contraindication for immunisation)
<p>Cautions including any relevant action to be taken</p> <p>Continued over page</p>	<p>Facilities for management of anaphylaxis should be available at all vaccination sites (see Chapter 8 of the Green Book) and advice issued by the Resuscitation Council UK.</p> <p>Tip cap of the syringe may contain natural rubber latex. For latex allergies other than anaphylactic allergies (such as a history of contact allergy to latex gloves), vaccines supplied in vials or syringes that contain latex can be administered.</p> <p>Very premature infants (born ≤ 28 weeks of gestation) who are in hospital should have respiratory monitoring for 48-72 hours when given their first immunisation, particularly those with a previous history of respiratory immaturity. If the child has apnoea, bradycardia</p>

² Exclusion under this PGD does not necessarily mean the medication is contraindicated, but it would be outside its remit and another form of authorisation will be required

<p>Cautions including any relevant action to be taken (continued)</p>	<p>or desaturations after the first immunisation, the second immunisation should also be given in hospital, with respiratory monitoring for 48-72 hours.</p> <p>The immunogenicity of the vaccine could be reduced in individuals who are immunosuppressed and individuals with HIV. However, vaccination should proceed in accordance with national recommendations see Chapter 22).</p> <p>Syncope (fainting) can occur following, or even before, any vaccination especially in adolescents as a psychogenic response to the needle injection. This can be accompanied by several neurological signs such as transient visual disturbance, paraesthesia and tonic-clonic limb movements during recovery. It is important that procedures are in place to avoid injury from faints.</p>
<p>Action to be taken if the patient is excluded</p>	<p>If aged less than 8 weeks 4CMenB is not routinely indicated, advise the parent/carer when the infant can be vaccinated. If aged from 2 years and not in a clinical risk group or requiring vaccination for the prevention of secondary cases of MenB disease, the individual/parent/carer should be advised that 4CMenB is not indicated. Individuals at increased risk of invasive meningococcal infection with asplenia, splenic dysfunction or complement disorders (including those on complement inhibitor treatment such as eculizumab) should be vaccinated in accordance with the recommended schedules in Chapter 7 and Chapter 22 of 'The Green Book' (see Meningococcal Group B Vaccine Risk Groups PGD).</p> <p>Individuals requiring vaccination for occupational health reasons should be referred to their occupational health service provider for vaccination.</p> <p>There are currently no recommendations for 4CMenB vaccination for individuals who are travelling or going to reside abroad.</p> <p>Individuals who have a history of anaphylactic allergy to latex should not be administered 4CMenB unless the benefit of vaccination outweighs the risk of an allergic reaction. Refer to appropriate clinician for assessment of risk: benefit – a PSD will be required.</p> <p>Individuals suffering acute severe febrile illness should postpone immunisation until they have recovered; immunisers should advise when the individual can be vaccinated and ensure another appointment is arranged.</p> <p>Seek appropriate advice from the local Screening and Immunisation Team, local Health Protection Team or the individual's clinician as required.</p> <p>The risk to the individual of not being immunised must be taken into account.</p> <p>Document the reason for exclusion and any action taken in the individual's clinical records.</p> <p>Inform or refer to the GP or a prescriber as appropriate.</p>
<p>Action to be taken if the patient or carer declines treatment (Continued over page)</p>	<p>Informed consent, from the individual or a person legally able to act on the person's behalf, must be obtained for each administration.</p> <p>Advise the individual/parent/carer about the protective effects of the vaccine, the risks of infection and potential complications of disease.</p>

Action to be taken if the patient or carer declines treatment (continued)	Document advice given and the decision reached. Inform or refer to the GP or a prescriber as appropriate.
Arrangements for referral for medical advice	As per local policy

Route and method of administration (continued)	matter and/or variation of physical aspect being observed, do not administer the vaccine. The vaccine's SPC provides further guidance on administration and is available from the electronic Medicines Compendium website .
Dose and frequency of administration	<p>Routine Immunisation Schedule</p> <p>The national recommendation for infants is a two dose primary course of 4CMenB, routinely starting at 8 weeks of age, to be administered with an 8 week interval and a booster dose to be administered, usually on or after their first birthday, although it may be administered until 2 years of age.</p> <p>4CMenB 0.5ml should ideally be given as follows:</p> <ul style="list-style-type: none"> • first primary immunisation visit (usually at age 8 weeks) • third primary immunisation visit (usually at age 16 weeks) • booster on or after the first birthday <p>Individuals with unknown or incomplete vaccination history</p> <p>Where there is no reliable history of previous immunisation, it should be assumed that they are unimmunised and the full UK recommendations should be followed (see Chapter 11).</p> <p>Infants younger than 12 months should receive the first dose of 4CMenB and second dose of 4CMenB two months later followed by the 4CMenB booster. Ensure that there is at least a two-month interval between the 4CMenB doses.</p> <p>Children aged one year to less than two years who received less than 2 4CMenB doses in the first year of life should receive two additional doses of 4CMenB at least two months apart.</p> <p>For further information see Guidance Vaccination of individuals with uncertain or incomplete immunisation status.</p> <p>Prevention of secondary cases of Men B disease</p> <p>Vaccination for the prevention of secondary cases of MenB disease should be given in accordance with recommendations from the UKHSA Health Protection Team and informed by the Guidance for Public Health Management of Meningococcal Disease in the UK.</p>
Duration of treatment	See dose section above
Quantity to be supplied and administered	Single dose of 0.5ml per an administration
Supplies	<p>Centrally purchased vaccines for the national immunisation programme can only be ordered via ImmForm. Vaccines for use for the national immunisation programme or for the prevention of secondary cases of MenB are provided free of charge.</p> <p>Vaccines for private prescriptions, occupational health use or travel or for individuals going to reside abroad are NOT provided free of charge and should be ordered from the manufacturer or wholesalers.</p> <p>Protocols for the ordering, storage and handling of vaccines should be followed to prevent vaccine wastage (see Green Book Chapter 3).</p>

Identification and management of adverse reactions (continued)	<p>4CMenB alone are similar to the other routine immunisations so paracetamol prophylaxis is not required.</p> <p>A detailed list of adverse reactions is available in the vaccine's SPC, which is available from the electronic Medicines Compendium website</p>
Reporting procedure of adverse reactions	<p>As with all vaccines, healthcare professionals and individuals/parents/carers are encouraged to report suspected adverse reactions to the Medicines and Healthcare products Regulatory Agency (MHRA) using the Yellow Card reporting scheme or search for MHRA Yellow Card in the Google Play or Apple App Store.</p> <p>Any adverse reaction to a vaccine should be documented in the individual's record and the individual's clinician should be informed.</p>
Written information to be given to patient or carer	<p>Offer marketing authorisation holder's patient information leaflet (PIL) provided with the vaccine.</p> <p>Immunisation promotional material may be provided as appropriate:</p> <ul style="list-style-type: none"> • Documents relating to the Meningococcal B (MenB) vaccination programme. • Protecting your baby against meningitis and septicaemia caused by meningococcal B bacteria • A guide to immunisations for babies up to 13 months of age • A quick guide to childhood immunisation for the parents of premature babies • Using paracetamol to prevent and treat fever after MenB vaccination (translated leaflets are also available to download from the health publications website) <p>Available from: www.gov.uk/government/collections/immunisation</p>
Patient advice and follow up treatment	<p>4CMenB is not expected to provide protection against all circulating meningococcal group B strains. Individuals should continue to seek prompt medical attention at the first signs of possible meningitis or septicaemia.</p> <p>Inform individuals who are immunosuppressed or individuals with HIV that the immunogenicity of the vaccine could be reduced.</p> <p>Inform individual/parent/carer of possible side effects and their management.</p> <p>If appropriate, advise the individual/parent/carer about the use and timing of paracetamol doses to reduce the risk, intensity and duration of fever (see Identification and management of adverse reactions).</p> <p>The individual/parent/carer should be advised to seek medical advice in the event of an adverse reaction or if they are concerned that their child is unwell at any time.</p> <p>When applicable, advise the individual/parent/carer when the subsequent vaccine dose is due.</p> <p>When administration is postponed advise the individual/parent/carer when to return for vaccination.</p>
Special considerations and additional information Continued over page	<p>It is important that premature infants have their immunisations at the appropriate chronological age, according to the schedule. . As the benefit of vaccination is high in premature and very premature infants, vaccination should not be withheld or delayed. The occurrence of apnoea following vaccination is especially increased in infants who were born very prematurely (see Cautions).</p>

<p>Special considerations and additional information (continued)</p>	<p>Meningococcal vaccines may be given to pregnant women when clinically indicated. There is no evidence of risk from vaccinating pregnant women or those who are breast-feeding with inactivated bacterial vaccines.</p> <p>Immunosuppression and HIV infection</p> <p>Individuals with immunosuppression and human immunodeficiency virus (HIV) infection (regardless of CD4 count) should be given meningococcal vaccines in accordance with the routine schedule (see Cautions).</p> <p>For further information on preventing secondary cases see the Public Health England Guidance for Public Health Management of Meningococcal Disease in the UK.</p>
<p>Records</p>	<p>Record:</p> <ul style="list-style-type: none"> • that valid informed consent was given • name of individual, address, date of birth and GP with whom the individual is registered • name of immuniser • name and brand of vaccine • date of administration • dose, form and route of administration of vaccine • quantity administered • batch number and expiry date • anatomical site of vaccination • advice given, including advice given if excluded or declines immunisation • details of any adverse drug reactions and actions taken • supplied via PGD <p>Records should be signed and dated (or a password controlled immuniser's record on e-records).</p> <p>All records should be clear, legible and contemporaneous.</p> <p>This information should be recorded in the individual's GP record. Where vaccine is administered outside the GP setting appropriate health records should be kept and the individual's GP informed.</p> <p>The local Child Health Information Systems team (Child Health Records Department) must be notified using the appropriate documentation/pathway as required by any local or contractual arrangement.</p> <p>A record of all individuals receiving treatment under this PGD should also be kept for audit purposes in accordance with local policy.</p>

6. Key references

Key references	Meningococcal B Vaccination <ul style="list-style-type: none">• Immunisation Against Infectious Disease: The Green Book, Chapter 4, last updated 20 March 2013, Chapter 7, last updated 10 January 2020 and Chapter 22 last updated 17 May 2022 www.gov.uk/government/collections/immunisation-against-infectious-disease-the-green-book• Bexsero® Summary of Product Characteristics, GlaxoSmithKline UK. Updated 19 June 2022. Bexsero Meningococcal Group B vaccine for injection in pre-filled syringe - Summary of Product Characteristics (SmPC) - (emc) (medicines.org.uk)• Meningococcal B (MenB) vaccination programme. Last updated 12 July 2021. www.gov.uk/government/collections/meningococcal-b-menb-vaccination-programme• Guidance for Public Health Management of Meningococcal Disease in the UK, Public Health England, updated 6 August 2019. www.gov.uk/government/publications/meningococcal-disease-guidance-on-public-health-management• Vaccination of individuals with uncertain or incomplete immunisation status. Public Health England. Updated 16 December 2019. www.gov.uk/government/publications/vaccination-of-individuals-with-uncertain-or-incomplete-immunisation-status• Meningococcal B: vaccine information for healthcare professionals 1 July 2021 https://www.gov.uk/government/publications/meningococcal-b-vaccine-information-for-healthcare-professionals General <ul style="list-style-type: none">• Health Technical Memorandum 07-01: Safe Management of Healthcare Waste. Department of Health 20 March 2013 www.england.nhs.uk/publication/management-and-disposal-of-healthcare-waste-htm-07-01/• National Minimum Standards and Core Curriculum for Immunisation Training. Published February 2018 www.gov.uk/government/publications/national-minimum-standards-and-core-curriculum-for-immunisation-training-for-registered-healthcare-practitioners• NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017. www.nice.org.uk/guidance/mpg2• NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 www.nice.org.uk/guidance/mpg2/resources• UKHSA Immunisation Collection www.gov.uk/government/collections/immunisation• Vaccine Incident Guidance www.gov.uk/government/publications/vaccine-incident-guidance-responding-to-vaccine-errors
-----------------------	---

7. Practitioner authorisation sheet

MenB PGD v6.00 Valid from: 28 February 2023 Expiry: 28 February 2025

Before signing this patient group direction (PGD), check that the document has had the necessary authorisations in section two. Without these, this PGD is not lawfully valid.

Practitioner

By signing this PGD you are indicating that you agree to its contents and that you will work within it.

PGDs do not remove inherent professional obligations or accountability.

It is the responsibility of each professional to practise only within the bounds of their own competence and professional code of conduct.

I confirm that I have read and understood the content of this PGD and that I am willing and competent to work to it within my professional code of conduct.			
Name	Designation	Signature	Date

Authorising manager

I confirm that the practitioners named above have declared themselves suitably trained and competent to work under this PGD. I give authorisation on behalf of <i>insert name of organisation</i> for the above named health care professionals who have signed the PGD to work under it.			
Name	Designation	Signature	Date

Note to authorising manager

Score through unused rows in the list of practitioners to prevent practitioner additions post managerial authorisation.

This authorisation sheet should be retained to serve as a record of those practitioners authorised to work under this PGD.