



PHE publications gateway number: GOV-9518

Zostavax[®] vaccine Patient Group Direction (PGD)

This PGD is for the administration of Zostavax[®], shingles (herpes zoster, live) vaccine, to individuals who are eligible for the national shingles immunisation programme for the prevention of herpes zoster ('zoster' or shingles) and herpes zoster-related post-herpetic neuralgia (PHN).

This PGD is for the administration of Zostavax[®] by registered healthcare practitioners identified in [Section 3](#), subject to any limitations to authorisation detailed in [Section 2](#).

Reference no: Zostavax PGD
Version no: v10.00
Valid from: 1 September 2021
Review date: 1 March 2023
Expiry date: 31 August 2023

Public Health England has developed this PGD to facilitate the delivery of publicly funded immunisation in line with national recommendations.

Those using this PGD must ensure that it is organisationally authorised and signed in Section 2 by an appropriate authorising person, relating to the class of person by whom the product is to be supplied, in accordance with Human Medicines Regulations 2012 (HMR2012)¹. **The PGD is not legal or valid without signed authorisation in accordance with [HMR2012 Schedule 16 Part 2](#).**

Authorising organisations must not alter, amend or add to the clinical content of this document (sections 4, 5 and 6); such action will invalidate the clinical sign-off with which it is provided. In addition, authorising organisations must not alter section 3 'Characteristics of staff'. Only sections 2 and 7 can be amended within the designated editable fields provided.

Operation of this PGD is the responsibility of commissioners and service providers. The final authorised copy of this PGD should be kept by the authorising organisation completing Section 2 for 8 years after the PGD expires as the PGD relates to adults only. Provider organisations adopting authorised versions of this PGD should also retain copies for 8 years.

Individual practitioners must be authorised by name, under the current version of this PGD before working according to it.

Practitioners and organisations must check that they are using the current version of the PGD. Amendments may become necessary prior to the published expiry date. Current versions of PHE PGD templates for authorisation can be found from: [Immunisation patient group direction \(PGD\) templates](#)

Any concerns regarding the content of this PGD should be addressed to: immunisation@phe.gov.uk

Enquiries relating to the availability of organisationally authorised PGDs and subsequent versions of this PGD should be directed to: england.swscreeningandimms@nhs.net.




¹ This includes any relevant amendments to legislation (such as [2013 No.235](#), [2015 No.178](#) and [2015 No.323](#)).
Zostavax PGD v10.00 Valid from: 01/09/2021 Expiry: 31/08/2023

Change history

Version number	Change details	Date
Final – revised 27 Aug 2013	New PHE PGD	Valid from 1 Sept 2013
Version 02.00	See earlier version of this PGD for change details.	4 June 2015
Version 03.00	See earlier version of this PGD for change details.	16 Nov 2015
Version 04.00	See earlier version of this PGD for change details.	3 Feb 2016
Version 05.00	See earlier version of this PGD for change details.	2 August 2016
Version 06.00	See earlier version of this PGD for change details.	07 April 2017
Version 07.00	See earlier version of this PGD for change details.	12 July 2017
Version 08.00	<p>PHE Shingles PGD amended to:</p> <ul style="list-style-type: none"> • include additional healthcare practitioners in Section 3 • refer to PHE Vaccine Incident Guidance in the off-label and storage sections • move the exclusion following natural infection to the cautions section and refer to the 'Shingles vaccination: Guidance for healthcare professionals' • include additional information in relation to the possible future availability of inactivated shingles vaccine • update off-label status section • include minor rewording, layout and formatting changes for clarity and consistency with other PHE PGD templates 	31 January 2019
Version 09.00	<p>PHE Shingles PGD amended to:</p> <ul style="list-style-type: none"> • reword inclusion criteria to remove catch-up cohort and define eligibility at 70 years and retention of eligibility until individuals 80th birthday • identify examples of biological therapy that are immunosuppressive monoclonal antibodies • include minor rewording, layout and formatting changes for clarity and consistency with other PHE PGD templates 	18 February 2021
Version 10.00	<p>PHE Shingles PGD amended to:</p> <ul style="list-style-type: none"> • rename as the Zostavax PGD and replace 'shingles (herpes zoster, live) vaccine' with 'Zostavax[®]' • reflect recommendations in the revised Green Book Chapter 28a and changes to the national shingles programme following the introduction of Shingrix[®] vaccine (see PHE Shingrix PGD) with amendment to the criteria for exclusion, cautions, actions if excluded, drug interaction and additional information sections • include minor rewording, layout and formatting changes for clarity and consistency with other PHE PGD templates and the Green Book Chapter 28a 	22 August 2021

1. PGD development

This PGD has been developed by the following health professionals on behalf of Public Health England:

Developed by:	Name	Signature	Date
Pharmacist (Lead Author)	Elizabeth Graham Lead Pharmacist, Immunisation and Countermeasures, PHE		25/08/2021
Doctor	Mary Ramsay Consultant Epidemiologist and Head of Immunisation and Countermeasures, PHE		25/08/2021
Registered Nurse (Chair of Expert Panel)	David Green Nurse Consultant, Immunisations and Countermeasures, PHE		25/08/2021

This PGD has been peer reviewed by the PHE Immunisations PGD Expert Panel in accordance with PHE PGD Policy. It has been ratified by the PHE Medicines Governance Group and the PHE Quality and Clinical Governance Delivery Board.

Expert Panel

Name	Designation
Nicholas Aigbogun	Consultant in Communicable Disease Control, Yorkshire and Humber Health Protection Team, Public Health England
Gayatri Amirthalingam	Consultant Epidemiologist, Public Health England
Sarah Dermont	Clinical Project Coordinator and Registered Midwife, NHS Infectious Diseases in Pregnancy Screening Programme, Public Health England
Ed Gardner	Advanced Paramedic Practitioner / Emergency Care Practitioner, Medicines Manager, Proactive Care Lead
Michelle Jones	Senior Medicines Optimisation Pharmacist, NHS Bristol North Somerset and South Gloucestershire CCG
Jacqueline Lamberty	Lead Pharmacist Medicines Management Services, Public Health England
Vanessa MacGregor	Consultant in Communicable Disease Control, Public Health England, East Midlands Health Protection Team
Alison Mackenzie	Consultant in Public Health Medicine / Screening and Immunisation Lead, Public Health England (South West) / NHS England and NHS Improvement South (South West)
Gill Marsh	Principal Screening and Immunisation Manager, Public Health England / NHS England and NHS Improvement (North West)
Lesley McFarlane	Screening and Immunisation Manager: Clinical (COVID-19 and Influenza), Public Health England / NHS England and NHS Improvement (Midlands)
Tushar Shah	Lead Pharmacy Advisor, NHS England and NHS Improvement (London Region)


2. Organisational authorisations

This PGD is not legally valid until it has had the relevant organisational authorisation.

It is the responsibility of the organisation that has legal authority to authorise the PGD, to ensure that all legal and governance requirements are met. The authorising body accepts governance responsibility for the appropriate use of the PGD.

NHS England & NHS Improvement (South West) authorises this PGD for use by the services or providers listed below:

Authorised for use by the following organisations and/or services
All NHS England & NHS Improvement commissioned immunisation services within <ul style="list-style-type: none"> • Bath & North East Somerset, Swindon, and Wiltshire • Bristol, North Somerset, and South Gloucestershire • Cornwall and the Isles of Scilly • Devon • Dorset • Gloucestershire • Somerset
Limitations to authorisation
This patient group direction (PGD) must only be used by the registered healthcare practitioners identified in Section 3 who have been named by their organisation to practice under it. The most recent in-date final version authorised by NHS England and NHS Improvement (South West) must be used.
This PGD includes vaccination of individuals across the national immunisation programme. Users of this PGD should note that where they are commissioned to immunise certain groups this PGD does not constitute permission to offer immunisation beyond the groups they are commissioned to immunise

Organisational approval (legal requirement)			
Role	Name	Sign	Date
Medical Director, System Improvement and Professional Standards, NHS England and NHS Improvement (South West)	Dr Kheelna Bavalia MRCGP MSc		1 September 2021

Additional signatories according to locally agreed policy			
Role	Name	Sign	Date

Local enquiries regarding the use of this PGD may be directed to england.swscreeningandimms@nhs.net.

Section 7 provides a practitioner authorisation sheet. Individual practitioners must be authorised by name to work to this PGD. Alternative practitioner authorisation sheets may be used where appropriate in accordance with local policy but this should be an individual agreement or a multiple practitioner authorisation sheet as included at the end of this PGD.

3. Characteristics of staff

<p>Qualifications and professional registration</p>	<p>Registered professional with one of the following bodies:</p> <ul style="list-style-type: none"> • nurses and midwives currently registered with the Nursing and Midwifery Council (NMC) • pharmacists currently registered with the General Pharmaceutical Council (GPhC) (Note: This PGD is not relevant to privately provided community pharmacy services) • paramedics and physiotherapists currently registered with Health and Care Professions Council (HCPC) <p>The practitioners above must also fulfil the Additional requirements detailed below.</p> <p>Check Section 2 Limitations to authorisation to confirm whether all practitioners listed above have organisational authorisation to work under this PGD.</p>
<p>Additional requirements</p>	<p>Additionally, practitioners:</p> <ul style="list-style-type: none"> • must be authorised by name as an approved practitioner under the current terms of this PGD before working to it • must have undertaken appropriate training for working under PGDs for supply/administration of medicines • must be competent in the use of PGDs (see NICE Competency framework for health professionals using PGDs) • must be familiar with the vaccine product and alert to changes in the Summary of Product Characteristics (SPC), Immunisation Against Infectious Disease ('The Green Book'), and national and local immunisation programmes • must have undertaken training appropriate to this PGD as required by local policy and in line with the National Minimum Standards and Core Curriculum for Immunisation Training • must be competent to undertake immunisation and to discuss issues related to immunisation • must be competent in the handling and storage of vaccines, and management of the 'cold chain' • must be competent in the recognition and management of anaphylaxis • must have access to the PGD and associated online resources • should fulfil any additional requirements defined by local policy <p>The individual practitioner must be authorised by name, under the current version of this PGD before working according to it.</p>
<p>Continued training requirements</p>	<p>Practitioners must ensure they are up to date with relevant issues and clinical skills relating to immunisation and management of anaphylaxis, with evidence of appropriate Continued Professional Development (CPD).</p> <p>Practitioners should be constantly alert to any subsequent recommendations from Public Health England and/or NHS England and NHS Improvement and other sources of medicines information. Note: The most current national recommendations should be followed but a Patient Specific Direction (PSD) may be required to administer the vaccine in line with updated recommendations that are outside the criteria specified in this PGD.</p>

4. Clinical condition or situation to which this PGD applies

<p>Clinical condition or situation to which this PGD applies</p>	<p>Zostavax® is indicated for vaccination of adults who are eligible for the national shingles immunisation programme for the prevention of herpes zoster ('zoster' or shingles) and herpes zoster-related post-herpetic neuralgia (PHN) in accordance with the recommendations given in Chapter 28a of Immunisation Against Infectious Disease: 'The Green Book'.</p>
<p>Criteria for inclusion</p>	<p>Individuals who:</p> <ul style="list-style-type: none"> • are aged 70 years (routine cohort) • have existing eligibility for Zostavax® under the national immunisation programme but remain unimmunised. Individuals from 70 years of age remain eligible for shingles immunisation until their 80th birthday.
<p>Criteria for exclusion²</p>	<p>Individuals for whom no valid consent has been received</p> <p>Individuals who:</p> <ul style="list-style-type: none"> • are under 70 years of age • are 80 years of age or over, even if they were previously in an eligible cohort • have had a confirmed anaphylactic reaction to a previous dose of varicella vaccine or to any component of Zostavax®, including neomycin or gelatin • are suffering from acute severe febrile illness (the presence of a minor infection is not a contraindication for immunisation) • have active untreated tuberculosis • have shingles infection with active lesions • have received systemic therapy in the last 48 hours with anti-viral medicines known to be effective against varicella zoster virus, such as aciclovir, • have received MMR vaccine in the preceding 4 weeks • are pregnant • are within 14 days of commencement of immunosuppressive therapy • have a primary or acquired immunodeficiency state as defined in Chapter 28a as a contraindication to Zostavax® administration • are on immunosuppressive or immunomodulating therapy as defined in Chapter 28a as a contraindication to Zostavax® administration <p>Note: Zostavax® is not contraindicated for use in individuals who are receiving topical or inhaled corticosteroids or corticosteroid replacement therapy. Individuals with low levels of immunosuppression can receive Zostavax®. If primary healthcare professionals administering the vaccine have concerns about the nature of therapies or the degree of immunosuppression, they should contact the relevant specialist for advice. Specialist advice should also be considered for individuals on combination immunosuppressive therapy.</p>

² Exclusion under this PGD does not necessarily mean the medication is contraindicated, but it would be outside its remit and another form of authorisation will be required
 Zostavax PGD v10.00 Valid from: 01/09/2021 Expiry: 31/08/2023

<p>Cautions including any relevant action to be taken</p>	<p>The decision to administer shingles vaccine to immunosuppressed individuals should be based on a clinical risk assessment. If the individual is under specialist care, and it is not possible to obtain full information on that individual's treatment history, then vaccination should not proceed until the advice of the specialist or a local immunologist has been sought.</p> <p>Individuals who have previously received immunosuppressive therapy should be carefully evaluated for the reconstitution of the immune system prior to receiving Zostavax®.</p> <p>Transmission</p> <p>There is a theoretical risk, in those who develop a rash following Zostavax® vaccination, of transmitting the attenuated vaccine virus to a susceptible individual. This risk should be weighed against the reduced risk of developing natural shingles and much higher risk of transmission from the circulating wild type varicella zoster virus in the community.</p> <p>As a precautionary measure, individuals who develop a vesicular rash after receiving Zostavax® should ensure the rash area is kept covered when in contact with a susceptible (chicken pox naïve) person until the rash is dry and crusted. If the individual with the vesicular rash are themselves immunosuppressed, they should avoid contact with susceptible people until the rash is dry and crusted, due to the higher risk of virus shedding. Prophylactic aciclovir can be considered in vulnerable patients exposed to a varicella like rash in a recent vaccinee.</p> <p>In the event of a person developing a varicella (widespread) or shingles-like (dermatomal) rash post-Zostavax®, a vesicle fluid sample should be sent for analysis, to confirm the diagnosis and determine whether the rash is vaccine-associated or wild-type. See Chapter 28a for more details.</p>
<p>Action to be taken if the patient is excluded</p> <p>Continued over page</p>	<p>Individuals who are not of eligible age for the national shingles immunisation programme should be advised when they will become eligible or why they are not eligible for immunisation.</p> <p>For individuals who have had a confirmed anaphylactic reaction to a previous dose of varicella vaccine or to any component of Zostavax® there is as an alternative inactivated vaccine Shingrix®. However, supply of Shingrix® is currently, at the time of writing, limited such that under the national shingles vaccination programme it is only recommended to be offered to immunosuppressed individuals (see Chapter 28a for current recommendations).</p> <p>Individuals suffering acute severe febrile illness should postpone immunisation until they have recovered.</p> <p>Individuals with untreated tuberculosis should postpone immunisation until their tuberculosis has been treated.</p> <p>Zostavax® is not recommended for the treatment of shingles. Individuals who have shingles should wait until symptoms have ceased before being considered for shingles immunisation. The natural boosting that occurs following an episode of shingles, however, makes the benefit of offering Zostavax® immediately following recovery unclear.</p>

<p>Action to be taken if the patient is excluded (continued)</p>	<p>Individuals who have received systemic anti-viral medicines known to be effective against varicella zoster virus, such as aciclovir, should postpone Zostavax[®] vaccination until at least 48 hours after cessation of treatment, as these medicines may reduce the response to the vaccine. The use of topical aciclovir is not a contraindication to vaccination.</p> <p>Individuals who have received MMR vaccine should postpone Zostavax[®] administration until a four-week minimum interval period has been observed.</p> <p>Individuals within 14 days of commencement of immunosuppressive therapy are not currently, at the time of writing, eligible for pre-treatment vaccination with Shingrix[®]. Supply of Shingrix[®] is currently, at the time of writing, limited and so vaccine supplied via the national programme should not be used for this indication. Eligible individuals who have not received Zostavax[®] are recommended to receive a single dose of vaccine at the earliest opportunity and at least 14 days before starting immunosuppressive therapy, although leaving one month would be preferable if a delay is possible (see Chapter 28a for current recommendations).</p> <p>Immunosuppressed individuals who are eligible for shingles vaccination but who are contraindicated to the receipt of the live vaccine Zostavax[®], in accordance with Chapter 28a should be offered Shingrix[®] instead (see PHE Shingrix PGD). If there is any doubt, individual patients should be discussed with their specialist.</p> <p>The risk to the individual of not being vaccinated must be taken into account.</p> <p>When administration is postponed arrange a future date for vaccination as appropriate, with due consideration of the individual's age to ensure they will meet the inclusion criteria for immunisation. If vaccination cannot be given before the individual is 80 years old explain why vaccination will no longer be indicated.</p> <p>Document the reason for exclusion and any action taken in individual's clinical records.</p> <p>Inform or refer to the GP or a prescriber as appropriate.</p>
<p>Action to be taken if the patient or carer declines treatment</p>	<p>Informed consent, from the individual or a person legally able to act on the individual's behalf, must be obtained prior to administration.</p> <p>Advise the individual/carer about the protective effects of the vaccine, the risks of infection and potential complications.</p> <p>Document advice given and the decision reached.</p> <p>Inform or refer to the GP or a prescriber as appropriate.</p>
<p>Arrangements for referral for medical advice</p>	<p>As per local policy</p>

5. Description of treatment

Name, strength and formulation of drug	<p>Zostavax[®], shingles (herpes zoster, live) vaccine, powder and solvent for suspension for injection.</p> <p>After reconstitution, Zostavax[®] lyophilised suspension (0.65ml) contains shingles (herpes zoster) vaccine, consisting of live attenuated virus derived from varicella zoster virus.</p>
Legal category	<p>Prescription only medicine (POM).</p>
Black triangle ▼	<p>No.</p>
Off-label use	<p>Vaccine should be stored according to the conditions detailed in the Storage section below. However, in the event of an inadvertent or unavoidable deviation of these conditions refer to PHE Vaccine Incident Guidance. Where vaccine is assessed in accordance with these guidelines as appropriate for continued use this would constitute off-label administration under this PGD.</p> <p>Where a vaccine is recommended off-label consider, as part of the consent process, informing the individual/carer that the vaccine is being offered in accordance with national guidance but that this is outside the product licence.</p>
Route / method of administration	<p>Following reconstitution, Zostavax[®] is given as a single dose by intramuscular or subcutaneous injection, preferably in the deltoid region of the upper arm. Intramuscular administration is preferred as injection-site adverse reactions were significantly less frequent in those who received the vaccine via this route.</p> <p>For individuals with a bleeding disorder, Zostavax[®] should be given by deep subcutaneous injection to reduce the risk of bleeding.</p> <p>Zostavax[®] should NOT be injected intravascularly.</p> <p>When administering at the same time as other vaccines care should be taken to ensure that the appropriate route of injection is used for all the vaccinations.</p> <p>The vaccines should be given at separate sites, preferably in different limbs. If given in the same limb, they should be given at least 2.5cm apart. The site at which each vaccine was given should be noted in the individual's records.</p> <p>It is recommended that the vaccine be administered immediately after reconstitution, to minimise loss of potency. Discard reconstituted vaccine if it is not used within 30 minutes.</p> <p>Avoid contact with disinfectants.</p> <p>When reconstituted, Zostavax[®] is a semi-hazy to translucent, off-white to pale yellow liquid. Discard the vaccine if there is any foreign particulate matter present or the appearance of the reconstituted vaccine differs from this description.</p> <p>The SPC for the vaccine provides further guidance on reconstitution and administration.</p>
Dose and frequency of administration	<p>Single dose of 0.65ml of reconstituted Zostavax[®].</p>

Duration of treatment	Single dose
Quantity to be supplied / administered	Single 0.65ml dose of reconstituted Zostavax®.
Supplies	<p>Centrally purchased vaccines for the national immunisation programme can only be ordered via ImmForm and are provided free of charge.</p> <p>Protocols for the ordering, storage and handling of vaccines should be followed to prevent vaccine wastage (see Green Book Chapter 3).</p>
Storage	<p>Store between +2°C to +8°C. Store in original packaging in order to protect from light. Do not freeze. Avoid contact with disinfectants.</p> <p>After reconstitution the vaccine should be used immediately. However, the in-use stability has been demonstrated for 30 minutes when stored at +20°C to +25°C.</p> <p>In the event of an inadvertent or unavoidable deviation of these conditions, vaccine that has been stored outside the conditions stated above should be quarantined and risk assessed for suitability of continued off-label use or appropriate disposal. Refer to PHE Vaccine Incident Guidance.</p>
Disposal	<p>Equipment used for immunisation, including used vials, ampoules, or discharged vaccines in a syringe or applicator, should be disposed of safely in a UN-approved puncture-resistant 'sharps' box, according to local authority arrangements and guidance in the technical memorandum 07-01 (Department of Health, 2013).</p>
Drug interactions	<p>None reported.</p> <p>See the Additional information section and SPC for information on co-administration with anti-virals and other vaccines.</p>
Identification and management of adverse reactions	<p>The most common adverse reactions observed after administration of Zostavax® are injection site reactions, including redness, swelling, pain and itching. Other relatively common reactions include bruising, hardening (induration) and warmth at the injection site, headache and pain in the relevant limb. Very rarely a varicella (chickenpox) - like illness has been reported.</p> <p>In the event of a person developing a varicella (widespread) or shingles-like (dermatomal) rash post-Zostavax® vaccination, a vesicle fluid sample should be sent for analysis, to confirm the diagnosis and determine whether the rash is vaccine-associated or wild-type (see Chapter 28a).</p> <p>A detailed list of adverse reactions is available in the SPC.</p>

<p>Special considerations / additional information continued</p>	<p>Immunisation with Zostavax® should ideally be delayed for seven days after COVID-19 vaccination and vice versa. Neither vaccine has been tested for routine co-administration; there may be a reduced response to Zostavax®. Where individuals attend requiring both vaccines, however, and require rapid protection or are considered likely to be lost to follow up, co-administration may still be considered</p> <p>There is no data on concomitant use with anti-viral medications known to be effective against varicella zoster virus but it is likely that these will reduce the response to Zostavax® - see Criteria for exclusion.</p> <p>See Chapter 28a for more details.</p>
<p>Records</p>	<p>Record:</p> <ul style="list-style-type: none"> • that valid informed consent was given • name of individual, address, date of birth and GP with whom the individual is registered • name of immuniser • name and brand of vaccine • date of administration • dose, form and route of administration of vaccine • quantity administered • batch number and expiry date • anatomical site of vaccination • advice given, including advice given if excluded or declines immunisation • details of any adverse drug reactions and actions taken • supplied via PGD <p>Records should be signed and dated (or a password controlled immuniser's record on e-records).</p> <p>All records should be clear, legible and contemporaneous.</p> <p>This information should be recorded in the individual's GP record and any other appropriate medical records, such as care or nursing records.</p> <p>A record of all individuals receiving treatment under this PGD should also be kept for audit purposes in accordance with local policy.</p>

6. Key references

Key references	<p>Shingles</p> <ul style="list-style-type: none">• Zostavax® Summary of Product Characteristics. MSD Ltd. Updated 8 May 2021. https://www.medicines.org.uk/emc/product/6101• Immunisation Against Infectious Disease: The Green Book, Chapter 28a. Updated 23 August 2021. https://www.gov.uk/government/collections/immunisation-against-infectious-disease-the-green-book• Shingles: Guidance and Vaccination Programme. Updated 18 August 2021. https://www.gov.uk/government/collections/shingles-vaccination-programme• Shingles vaccination: Guidance for healthcare professionals. Public Health England. Published March 2018. https://www.gov.uk/government/publications/shingles-vaccination-guidance-for-healthcare-professionals <p>General</p> <ul style="list-style-type: none">• Health Technical Memorandum 07-01: Safe Management of Healthcare Waste. Department of Health. 20 March 2013. https://www.gov.uk/government/publications/guidance-on-the-safe-management-of-healthcare-waste• National Minimum Standards and Core Curriculum for Immunisation Training. Published February 2018. https://www.gov.uk/government/publications/national-minimum-standards-and-core-curriculum-for-immunisation-training-for-registered-healthcare-practitioners• NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017. https://www.nice.org.uk/guidance/mpg2• NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017. https://www.nice.org.uk/guidance/mpg2/resources• PHE Immunisation Collection. https://www.gov.uk/government/collections/immunisation• PHE Vaccine Incident Guidance. https://www.gov.uk/government/publications/vaccine-incident-guidance-responding-to-vaccine-errors
-----------------------	---

7. Practitioner authorisation sheet

Zostavax PGD v010.00 Valid from: 01/09/2021 Expiry: 31/08/2023

Before signing this patient group direction (PGD), check that the document has had the necessary authorisations in section two. Without these, this PGD is not lawfully valid.

Practitioner

By signing this PGD you are indicating that you agree to its contents and that you will work within it.

PGDs do not remove inherent professional obligations or accountability.

It is the responsibility of each professional to practise only within the bounds of their own competence and professional code of conduct.

I confirm that I have read and understood the content of this PGD and that I am willing and competent to work to it within my professional code of conduct.			
Name	Designation	Signature	Date

Authorising manager

I confirm that the practitioners named above have declared themselves suitably trained and competent to work under this PGD. I give authorisation on behalf of <i>insert name of organisation</i> for the above named health care professionals who have signed the PGD to work under it.			
Name	Designation	Signature	Date

Note to authorising manager

Score through unused rows in the list of practitioners to prevent practitioner additions post managerial authorisation.

This authorisation sheet should be retained to serve as a record of those practitioners authorised to work under this PGD.