



Department  
of Health



Public Health  
England

**NHS**  
England

NHS England Publications Gateway Reference: 00360

Dear Nurse/Midwife,

As we approach our busiest season, influenza outbreaks become a greater threat to ourselves and our patients. It is imperative that you take steps to protect those you care for, colleagues, family and yourself against flu by **getting vaccinated**.

Nurses, midwives and care staff form the largest staff body within the NHS and social care, and as such, we can make the biggest impact in preventing the spread of flu. As well as taking action to protect yourself, you have the potential to influence and encourage your colleagues to get their flu jab. Please use the attached evidence to dispel any misconceptions you or your colleagues may have, by reinforcing the benefits of immunisation and reassuring people that it is safe and effective.

By protecting yourself against flu, you not only set an example to colleagues and peers but also help to continue providing high quality compassionate care to those we care for.

For more information about the national seasonal flu campaign for staff visit <http://www.nhsemployers.org/flu>.

Thank you for your support.

A handwritten signature in black ink, appearing to read 'Jane Cummings'.

Jane Cummings  
Chief Nursing Officer  
NHS England

A handwritten signature in black ink, appearing to read 'Viv Bennett'.

Viv Bennett  
Director of Nursing,  
Public Health England  
and Department of Health

A handwritten signature in black ink, appearing to read 'D M Salisbury'.

Professor D M SALISBURY  
Director of Immunisation, Department of Health

# The importance of influenza vaccination for Nurses and Midwives

Every year influenza vaccination is offered to NHS staff as a way to reduce the risk of staff and patients contracting and transmitting the virus. Vaccine uptake across NHS organisations varies from below 10% to above 90% with a national uptake of 45.6% (2012/13). This document summarises the evidence showing why vaccination of doctors and healthcare workers is important and answers some common questions.

## Why should we worry about influenza?

Influenza can cause a spectrum of illness from mild to severe, even among people who are previously well. There were 457 confirmed deaths from influenza reported in 2009-10 and 602 in 2010-11<sup>ii</sup>. Nearly 9000 patients were admitted to hospital with influenza in England in 2010-11, of whom 2200 were admitted to intensive care<sup>iii</sup>. While the impact of influenza was less marked during the 2012-13 season, influenza remains unpredictable and it is hard to forecast the severity of future influenza seasons.

These figures are high for a disease that is largely preventable through vaccination. As a comparison, hepatitis B causes around 60 deaths per year.<sup>iv</sup> The hepatitis B vaccine is an immunisation that healthcare professionals in the NHS are expected to have. Influenza deaths are also high compared to other infectious diseases: invasive meningococcal disease causes around 60-80 deaths per year<sup>viii</sup>.

## Why is vaccination important for clinical staff?

### *Protecting yourself*

Frontline healthcare workers are more likely to be exposed to the influenza virus, particularly during winter months when some of their patients will be infected. It has been estimated that up to 1 in 4 healthcare workers may become infected with influenza during a mild influenza season, a much higher incidence than expected in the general population<sup>v</sup>.

Even previously healthy people and the young can develop severe complications from influenza; up to one third of deaths in 2009-10 and 2010-11 were in people considered healthy<sup>vi</sup>, with many of the cases of severe illness in those aged under 65 years<sup>ii, v, vii, viii, ix</sup> (89% of hospital admissions, 87% of critical care beds occupied and 79% of deaths).

### *Protecting your patients*

Influenza is a highly transmissible infection. The patient population found in hospital is much more vulnerable to the severe effects of influenza<sup>x</sup>. Healthcare workers may transmit the illness to patients even if they are mildly or sub-clinically infected. There are reports of influenza outbreaks within hospitals and other care settings where transmission from healthcare workers to patients is likely to have facilitated spread of the disease<sup>xi, xii, xiii</sup>. In an influenza outbreak in one trust 118 staff and 49 patients were infected<sup>xi</sup>. A second resulted in six infections among neonates and one death<sup>xiii</sup>.

'Herd-immunity' of healthcare workers to reduce the likelihood of introduction and transmission of the virus in care settings is an effective way to prevent this. Settings randomised to high levels

of immunisation had reduced rates of flu-like illness, hospitalisation and mortality in the elderly in comparison with controls<sup>xiv,xv,xvi,xvii</sup>.

*Protecting your friends, family and colleagues.*

Some healthcare workers, aware that they are more likely to become infected with influenza, get the flu vaccination in order to protect other family from influenza, particularly young children or other relatives who may fall into at-risk groups<sup>xviii</sup>.

### **Advice from professional bodies**

The Green Book recommends that healthcare workers directly involved in patient care be vaccinated annually<sup>x</sup>. It is also encouraged by the General Medical Council in section 29 of the good medical practice<sup>xix</sup>, and by the BMA<sup>xx</sup>.

### **How effective is the vaccine?**

A recent meta-analysis, which included studies when the influenza virus strains in the vaccine were drifted or mismatched with those in circulation, suggested an overall efficacy against confirmed disease of 59% (95% confidence interval 51-67) in adults aged 18 to 65 years<sup>xxi</sup>

### **How safe is the vaccine?**

The most common side effect is bruising or local muscular stiffness (10-64%) at the injection site<sup>xxii</sup>. Other reported side-effects following the vaccine include fever, malaise and myalgia. These are short lived and their incidence may not be much greater in comparison with those who receive a placebo vaccine (fever 3% vs 1%; malaise 9% vs 6%; myalgia 18% vs 10%)<sup>xxiii</sup>. Some of these side effects were particularly common during the pandemic, as the vaccines used then had an adjuvant. The present trivalent vaccine does not contain adjuvants so such side effects will be less common.

Although it is common for people to complain that the vaccine gave them influenza, this is not possible. It is most likely that flu-like symptoms experienced by people who have just had the vaccine are not caused by influenza but are the result of many other circulating viruses that can produce influenza-like symptoms. It also takes up to two weeks to develop immunity after vaccination, so infection could occur during this window.

### **What about severe reactions?**

The risk of having an anaphylactic reaction to the seasonal influenza vaccine is very rare, but those who have had a severe reaction (anaphylaxis) to a previous dose of seasonal influenza vaccine or to any part of the vaccine should not receive it. Individuals who have egg allergy may be at increased risk of reaction to influenza vaccines. In recent years, inactivated influenza vaccines that are egg-free or have a very low ovalbumin content have become available. Patients who have either confirmed anaphylaxis to egg or egg allergy with uncontrolled asthma (BTS SIGN step 4 or above) can be immunised with an egg-free influenza vaccine.

If no egg-free vaccine is available at the clinic, patients should be referred to specialists for vaccination in hospital using an inactivated influenza vaccine with an ovalbumin content less than 0.12 µg/ml. A split dose schedule may be required at the discretion of the supervising physician. Facilities should be available and staff trained to recognise and treat anaphylaxis. Vaccines with ovalbumin content more than 0.12 µg/ml or where content is not stated should not be used in egg-allergic individuals.

All other egg allergic individuals can be given egg-free vaccine or inactivated influenza vaccine with an ovalbumin content less than 0.12 µg/ml administered as recommended in primary care.

More detailed information on the characteristics of the available vaccines for 2012-13, including age indications and ovalbumin (egg) content can be found in the Seasonal Flu chapter of the Green Book.

### How is safety of the vaccines monitored?

As with all medicines in the UK, influenza vaccines require licensing by the Medicines and Healthcare Products Regulatory Agency (MHRA). Like other medical products, passive surveillance, using reports from yellow cards, is used to identify adverse events. The observed rate of adverse reports is compared to the expected rate, based on data from a general practice research database, after making allowance for under-reporting.

This is complemented by active surveillance, which uses very large population cohorts from primary care databases to proactively look at the risk of an adverse event which may be of concern. Comparisons are made between patterns of self-presenting illness to general practice in the period after vaccination compared to controls. Other countries have similar systems and data is pooled and reviewed at national and international levels.

### Why do some nurses or midwives refuse the vaccine?

There are a variety of reasons why staff decline the vaccine. A recent survey of healthcare workers in University Hospitals of Leicester and Leicestershire Partnership Trust<sup>xxiv</sup> found that one third of unvaccinated clinician respondents felt that universal infection control practices are sufficient. One third of unvaccinated clinician respondents reported they were not vaccinated because they have a good diet and/or take vitamins or supplements that work as well as or better than the influenza vaccine.

Vaccination is the best option for protecting yourself, your family and vulnerable patients from the virus. Although infection control measures are vital and a good diet is encouraged, neither action alone will prevent influenza.

### When should I be vaccinated?

The new vaccines should be available from the end of September 2013 and any healthcare worker with direct patient contact is urged to get vaccinated as soon as possible. Your local occupational health department is likely to lead on delivery so the advice is to contact them or the appropriate team in September. Any healthcare workers in at-risk groups can receive the vaccine at their GP, but are asked to please report this vaccination at work to ensure inclusion in uptake figures recorded for the Department of Health.

---

i Hinde D. The 2009 influenza pandemic – an independent review of the UK response to the 2009 pandemic. The Cabinet Office, London, 2010.

ii Health Protection Agency. Surveillance of influenza and other respiratory viruses in the UK: 2010-2011 report. HPA, London, 2011.

iii Estimates of hospital admissions from Hospital Episode Statistics 2010-2011; estimates of critical care admissions taken from HPA annual influenza report based on bed-days and a mean length of stay of seven days.

iv Office of National Statistics. Mortality Statistics: Deaths registered in England and Wales. ONS, London (based on data from 2006-2010).

- v Elder AG, O'Donnell B, McCruden EA, Symington IS, Carman WF. Incidence and recall of influenza in a cohort of Glasgow healthcare workers during the 1993-4 epidemic: results of serum testing and questionnaire. *The British Medical Journal*, 1996;313:1241-2.
- vi Donaldson L, Rutter P, Ellis B et al. Pandemic Flu Mortality in England. *The British Medical Journal*, 2009;339:b5213
- vii Health Protection Agency. Epidemiological report of the 2009 pandemic (H1N1) 2009 in the UK. HPA, London, 2011.
- viii Campbell CN, Mytton OT, McLean EM et al. Hospitalization in two waves of pandemic influenza A(H1N1) in England. *Epidemiology and Infection*. 2011;139(10):1560-9.
- ix Mytton OT, Rutter PD, Mak M et al. Mortality due to pandemic (H1N1) 2009 influenza in England: a comparison of the first and second waves. *Epidemiology and Infection*. Published online: 1 Nov 2011.
- x Salisbury D, Ramsay M, Noakes K. Immunisation against infectious disease – „the Green Book“. Department of Health, 2011.
- xi Pachucki CT, Pappas SA, Fuller GF et al. Influenza A among hospital personnel and patients. Implications for recognition, prevention and control. *Archives of Internal Medicine* 1989; 149:77-80.
- xii Horcajada JP, Pumarola T, Martinez JA et al. A nosocomial outbreak of influenza during a period without influenza epidemic activity. *European Respiratory Journal* 2003; 21:303-7.
- xiii Cunney RJ, Bialachowski A, Thornley D, Smaill FM, Pennie RA. An outbreak of influenza A in a neonatal intensive care unit. *Infection Control and Hospital Epidemiology*. 200; 21:449-54
- xiv Potter J, Stott DJ, Roberts MA et al. The influenza vaccination of healthcare workers in long-term care hospitals reduces the mortality of elderly patients. *Journal of Infectious Diseases*. 1997; 175:1-6.
- xv Carman WF, Elder AG, Wallace LA et al. Effects of influenza vaccination of healthcare workers on mortality of elderly people in long term care: a randomised control trial. *The Lancet* 2000; 355:93-97.
- xvi Hayward AV, Harling R, Wetten S et al. Effectiveness of an influenza vaccine programme for care home staff to prevent death, morbidity, and health service use among residents: cluster randomised controlled trials. *The British Medical Journal* 2006; doi:10.1136/bmj.39010.581354.55.
- xvii Lemaitre M, Meret T, Rothan-Tondeur M et al. Effect of influenza vaccination of nursing home staff on mortality of residents: a cluster randomised trial. *Journal of American Geriatric Society* 2009; 57:1580-6.
- xviii Hollymeyer HG, Hayden F, Poland G, Buchholz U. Influenza vaccination of healthcare workers in hospitals – a review of studies on attitudes and predictors. *Vaccine* 2009; 27: 3935-44.
- xix The General Medical Council. *Good Medical Practice*. The GMC, London, 2013.
- xx Buckman L, Porter M, Shanbhag R, Dolphin T, Datta S. A message for doctors. *BMA letter to doctors*, 2011. NHS Employers.
- xxi Osterholm MT, Kelley NS, Sommer A, Belongia EA. Efficacy and effectiveness of influenza vaccines: a systematic review and meta-analysis. *Lancet Infect Dis*, 2012; 12: 36-44..
- xxii Centres for Disease Control. 2010-11 Influenza Prevention and Control Recommendations – Adverse Events After Receipt of TIV. CDC, Atlanta, 2011: [www.cdc.gov/flu/professionals/acip/adversetiv.htm](http://www.cdc.gov/flu/professionals/acip/adversetiv.htm)
- xxiii Jackson LA, Gaglani MJ, Keyserling HL et al. Safety, efficacy and immunogenicity of an inactivated influenza vaccine in healthy adults: a randomised, placebo controlled trial over two influenza seasons. *BMC Infectious Disease*, 2010; 10:71. doi. 10.1186/1471-2334-10-71
- xxiv Burch, T. (2012) “Motivators and barriers for influenza vaccine uptake amongst healthcare workers: results of an online staff survey”, East Midlands Seasonal Influenza Debrief, Nottingham.