

National Dose Banding Table – Eribulin*

Vial Sizes of Drug	0.88mg/2mL 1.32mg/3mL	Drug	Eribulin
--------------------	--------------------------	------	----------

* All doses and ranges are quoted as eribulin (without salt)

0.88mg eribulin (without salt) = 1mg eribulin mesylate

See table usage notes below regarding 'single container' and 'multiple syringe' tables.

Master Bands and Ranges

This table is intended to be in a format useful for electronic prescribing systems.

Use To (A) if your system will round UP for doses on the step between two bands.

Use To (B) if your system will round DOWN for doses on the step between two bands.

Note on accuracy: All dose bands and range figures are rounded to 2 decimal places. Due to the concentration of the drug almost all figures resulting from calculations have six or more significant figures which lies well in excess of either measurability or any reasonable requirement of accuracy.

Band Range (mg)			Band Dose (mg)	Variance (percent)	
From ≥	To (A) <	To (B) ≤		Below	Above
0.68	0.73	0.72	0.70	3	-3
0.73	0.77	0.76	0.75	3	-3
0.77	0.84	0.83	0.79	3	-5
0.84	0.92	0.91	0.88	5	-5
0.92	1.01	1.00	0.97	5	-4
1.01	1.10	1.09	1.06	4	-4
1.10	1.19	1.18	1.14	4	-4
1.19	1.28	1.27	1.23	4	-3
1.28	1.36	1.35	1.32	4	-3
1.36	1.45	1.44	1.41	3	-3
1.45	1.56	1.55	1.50	3	-4
1.56	1.69	1.68	1.63	4	-4
1.69	1.83	1.82	1.76	4	-4
1.83	1.96	1.95	1.89	4	-3
1.96	2.11	2.10	2.02	3	-4
2.11	2.31	2.30	2.20	4	-5
2.31	2.53	2.52	2.42	5	-4
2.53	2.75	2.74	2.64	4	-4
2.75	2.97	2.96	2.86	4	-4
2.97	3.19	3.18	3.08	4	-3
3.19	3.41	3.40	3.30	4	-3
3.41	3.73	3.72	3.52	3	-6

Drug- and Dose-specific Tables

These tables are intended to assist with non-electronic systems where protocol-specific information is required. The range of the tables has been extended beyond 'normal' use, especially in the lower ranges, to allow for up to a 50% dose reduction to be used in patients who are $1.3m^2$. Doses which are normally capped are highlighted in grey.

Eribulin – in 100mL sodium chloride 0.9% (note doses quoted as eribulin, not eribulin mesylate)

BSA	Choose nearest for dose reductions							1.0	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0
1.23 mg/m ²	0.7	0.75	0.79	0.88	0.97	1.06	1.14	1.23	1.32	1.5	1.63	1.76	1.89	2.02	2.02	2.2	2.42	2.42	2.64	2.64	2.86	2.86	3.08	3.3	3.3	3.52	3.52	3.52
0.97 mg/m ²	0.57	0.62	0.66	0.7	0.75	0.79	0.88	0.97	1.06	1.14	1.23	1.32	1.5	1.5	1.63	1.76	1.89	1.89	2.02	2.2	2.2	2.42	2.42	2.42	2.64	2.64	2.86	2.86

Table usage notes: 'Single container' tables are for products which are able to be supplied as a single unit, such as an infusion bag, one syringe, or as an infusor device (pump). 'Multiple syringe' (e.g. 'pick-and-mix') tables are for products which are supplied in combinations of syringes to make up a single dose. Some drugs may have both a single container table, and a multiple syringe table. Doses may vary between the two tables due to rounding requirements.

