

Name of product	Cisplatin infusion. Aseptically prepared from licensed sterile starting materials.
Concentration	Concentration range between 0.02mg/ml to 0.5mg/ml in accordance with the nat dose banding tables, and in an infusion bag volume determined by the local cent hydration regimens. The Cisplatin concentrate should be added to the bag without withdrawal of equiv volume of Sodium Chloride 0.9% w/v Infusion unless withdrawal of some infusion required to accomodate the total dose.
Diluent	Sodium Chloride 0.9% w/v
Volume	Typically 250ml to 1000ml as determined by the local centre's hydration regimen
Final container	Non-PVC e.g. polyolefin infusion bag with additive port cover. Ideally infusion bag design will incorporate self sealing giving port to minimise Harisks associated with accidental spillage during administration.
Starting materials	Licensed Cisplatin Injection 1mg/ml Licensed Sodium Chloride 0.9% w/v Infusion bags
Labelling	Labelling must be compliant with the principles of labelling for safety and the BP General specification on unlicensed medicines. Tall Man lettering must be used for drug name.
Label sample	An example label is provided below stating the minimum requirements only (the format is not restrictive and suppliers can use their preferred layout):- CISplatin xxxmg in xxxxml Sodium Chloride 0.9% w/v For Intravenous Infusion Infuse the entire contents of the bag Store at Room Temperature Protect From Light Expiry: dd/mm/yyyy BN: XXXXXXXXX Keep out of the reach and sight of children Manufacturer's details MS XXXXXX Caution Cytotoxic: Handle with care
Batch Number	All products will have a unique batch identification number
Latex status of - components - manufacturing process	All materials and manufacturing processes will be latex free or clearly labelled if
Stability	Stability studies should conform to the Standard Protocol for deriving and assess of stability of Aseptic preparations (small molecules) published by the NHS Pharmaceutical QA Committee (4th Edition, April 2017).