

Name of product	Decification: Vinblastine 50ml infusion bags Vinblastine infusion. Aseptically prepared from licensed sterile starting material
Concentration	Concentration range 0.1mg/ml to 0.3mg/ml in 50ml infusion bags in accordance
	the national dose banding tables. The Vinblastine concentrate should be added to the bag without withdrawal of
	equivalent volume of Sodium Chloride 0.9% w/v Infusion unless withdrawal of so
	infusion fluid is required to accomodate the total dose.
Diluent	Sodium Chloride 0.9% w/v
Volume	50ml
Final container	Non-PVC e.g. polyolefin infusion bag with additive port cover.
	Ideally infusion bag design will incorporate self sealing giving port to minimise
	risks associated with accidental spillage during administration.
Starting materials	Licensed Vinblastine Injection
Claring materials	Licensed Sodium Chloride 0.9% w/v Infusion bags
Laballia e	
Labelling	Labelling must be compliant with the principles of labelling for safety (in particular comply with HSC 2008/001 Updated national guidance on the
	safe administration of intrathecal chemotherapy and NPSA/2008/RRR004 i.e. r
	labelled 'For Intravenous Use Only – Fatal If Given by Other Routes') and the B
	General specification on unlicensed medicines. Tall Man lettering must be used
	drug name.
Label sample	An example label is provided below stating the minimum requirements only (the format is not restrictive and suppliers can use their preferred layout):-
	vinBLAStine xxmg in xxml Sodium Chloride 0.9% w/v
	in xxml Sodium Chloride 0.9% w/v FOR INTRAVENOUS USE ONLY
	FATALIF GIVEN BY OTHER ROUTES
	Infuse the entire contents of the bag
	Ctore in a Defricerator at 2 000 Dratest From Light
	Store in a Refrigerator at 2-8°C Protect From Light Expiry: dd/mm/yyyy BN: XXXXXXXXX
	Keep out of the reach and sight of children
	Manufacturer's details MS XXXXXX Caution Cytotoxic: Handle with care
Batch Number	All products will have a unique batch identification number
Latex status of	All materials and manufacturing processes will be latex free or clearly labelled
- components	
- manufacturing process	Stability studies should conform to the Standard Protocol for deriving and asse
Stability	Graphing studies should conform to the Standard Flotocol for denting and asset
	of stability of Aseptic preparations (small molecules) published by the NHS