

| Name of product | Amsacrine infusion. Aseptically prepared from licensed sterile starting materials |
|--|--|
| Concentration | Concentration range typically up to 0.5mg/ml in 500ml infusion bags in accordant with the national dose banding tables. The Amsacrine concentrate should be added to the bag without withdrawal of equivalent volume of Glucose 5% w/v Infusion unless withdrawal of some infusion fluid is required to accomodate the total dose. |
| Diluent | Glucose 5% w/v |
| Volume | 500ml |
| Final container | Non-PVC e.g. polyolefin infusion bag with additive port cover. Ideally infusion bag design will incorporate self sealing giving port to minimise F risks associated with accidental spillage during administration. |
| Starting materials | Licensed Amsacrine 50mg/ml concentrate solution for injection (75mg in 1.5ml) Licensed Amsacrine diluent 13.5ml Licensed Glucose 5% w/v Infusion bags |
| Labelling | Labelling must be compliant with the principles of labelling for safety and the BF General specification on unlicensed medicines. There is no requirement for Tal lettering for the drug name. |
| | format is not restrictive and suppliers can use their preferred layout):- Amsacrine xxxmg in xxxml Glucose 5% w/v For Intravenous Infusion Infuse the entire contents of the bag Store at Room Temperature Protect From Light Expiry: dd/mm/yyyy BN: XXXXXXXX Keep out of the reach and sight of children Manufacturer's details MS XXXXXX Caution Cytotoxic: Handle with care . |
| Batch Number | All products will have a unique batch identification number |
| Latex status of - components - manufacturing process | All materials and manufacturing processes will be latex free or clearly labelled it |
| Stability | Stability studies should conform to the Standard Protocol for deriving and asses of stability of Aseptic preparations (small molecules) published by the NHS |