

Name of product	Arsenic Trioxide infusion. Aseptically prepared from licensed sterile starting mat
Concentration	Concentration range typically up to 0.4mg/ml in 100ml infusion bags in accordant with the national dose banding tables. The Arsenic Trioxide concentrate should be added to the bag without withdrawa equivalent volume of Sodium Chloride 0.9% w/v Infusion unless withdrawal of s infusion fluid is required to accomodate the total dose.
Diluent	Sodium Chloride 0.9% w/v
Volume	100ml
Final container	Non-PVC e.g. polyolefin infusion bag with additive port cover. Ideally infusion bag design will incorporate self sealing giving port to minimise H risks associated with accidental spillage during administration.
Starting materials	Licensed Arsenic Trioxide 1mg/ml solution for injection Licensed Sodium Chloride 0.9% w/v Infusion bags
Labelling	Labelling must be compliant with the principles of labelling for safety and the BF General specification on unlicensed medicines. There is no requirement for Tal lettering for the drug name.
Label sample	An example label is provided below stating the minimum requirements only (the format is not restrictive and suppliers can use their preferred layout):- Arsenic Trioxide xxmg in xxxml Sodium Chloride 0.9% w/v For Intravenous Infusion Infuse the entire contents of the bag Store in a Refrigerator at 2-8°C Protect From Light Expiry: dd/mm/yyyy BN: XXXXXXXX Keep out of the reach and sight of children Manufacturer's details MS XXXXXX Caution Cytotoxic: Handle with care
Batch Number	All products will have a unique batch identification number
Latex status of - components - manufacturing process	All materials and manufacturing processes will be latex free or clearly labelled if
Stability	Stability studies should conform to the Standard Protocol for deriving and asses of stability of Aseptic preparations (small molecules) published by the NHS Pharmaceutical QA Committee (4th Edition, April 2017).