

National Dose Banding Table – Single Container

Strength of raw material (after reconstitution if required)	2.5 mg/mL	Example drug(s)	Bendamustine Bortezomib SC
--	-----------	-----------------	-------------------------------

See table usage notes below regarding 'single container' and 'multiple syringe' tables.

Master Bands and Ranges

This table is intended to be in a format useful for electronic prescribing systems.

Use To (A) if your system will round UP for doses on the step between two bands.

Use To (B) if your system will round DOWN for doses on the step between two bands.

Band Range (mg)			Band Dose (mg) *	Variance (percent)	
From ≥	To (A) <	To (B) ≤		Below	Above
0.58	0.63	0.62	0.6	4	-4
0.63	0.68	0.67	0.65	4	-4
0.68	0.73	0.72	0.7	4	-3
0.73	0.80	0.79	0.75	4	-6
0.80	0.90	0.89	0.85	6	-5
0.90	1.00	0.99	0.95	6	-5
1.00	1.10	1.09	1.05	5	-4
1.10	1.22	1.21	1.15	5	-6
1.22	1.37	1.36	1.3	6	-5
1.37	1.52	1.51	1.45	6	-5
1.52	1.70	1.69	1.6	5	-6
1.70	1.90	1.89	1.8	6	-5
1.90	2.12	2.11	2	5	-6
2.12	2.37	2.36	2.25	6	-5
2.37	2.62	2.61	2.5	5	-5
2.62	2.87	2.86	2.75	5	-4
2.87	3.12	3.11	3	4	-4
3.12	3.37	3.36	3.25	4	-4
3.37	3.62	3.61	3.5	4	-3
3.62	3.87	3.86	3.75	4	-3
3.87	4.12	4.11	4	3	-3
4.12	4.37	4.36	4.25	3	-3
4.37	4.74	4.73	4.5	3	-5
4.74	5.25	5.24	5	5	-5
5.25	5.75	5.74	5.5	5	-4
5.75	6.25	6.24	6	4	-4
6.25	6.75	6.74	6.5	4	-4
6.75	7.25	7.24	7	4	-3
7.25	7.75	7.74	7.5	4	-3
7.75	8.25	8.24	8	3	-3
8.25	8.75	8.74	8.5	3	-3
8.75	9.49	9.48	9	3	-5
9.49	10.49	10.48	10	5	-5
10.49	11.49	11.48	11	5	-4
11.49	12.73	12.72	12	4	-6
12.73	14.23	14.22	13.5	6	-5
14.23	15.73	15.72	15	5	-5
15.73	17.47	17.46	16.5	5	-6
17.47	19.48	19.47	18.5	6	-5
19.48	21.48	21.47	20.5	5	-5
21.48	23.72	23.71	22.5	5	-5
23.72	26.22	26.21	25	5	-5



Band Range (mg)			Band Dose (mg) *	Variance (percent)	
From ≥	To (A) <	To (B) ≤		Below	Above
26.22	28.72	28.71	27.5	5	-4
28.72	31.23	31.22	30	4	-4
31.23	33.73	33.72	32.5	4	-4
33.73	36.23	36.22	35	4	-3
36.23	38.73	38.72	37.5	4	-3
38.73	42.43	42.42	40	3	-6
42.43	47.44	47.43	45	6	-5
47.44	52.44	52.43	50	5	-5
52.44	57.45	57.44	55	5	-4
57.45	63.64	63.63	60	4	-6
63.64	71.15	71.14	67.5	6	-5
71.15	78.66	78.65	75	5	-5
78.66	86.17	86.16	82.5	5	-4
86.17	94.87	94.86	90	4	-5
94.87	106.07	106.06	100	5	-6
106.07	118.59	118.58	112.5	6	-5
118.59	132.29	132.28	125	5	-6
132.29	148.49	148.48	140	6	-6
148.49	166.02	166.01	157.5	6	-5
166.02	185.91	185.90	175	5	-6
185.91	209.63	209.62	197.5	6	-6
209.63	235.85	235.84	222.5	6	-6
235.85	265.75	265.74	250	6	-6
265.75	300.67	300.66	282.5	6	-6
300.67	340.59	340.58	320	6	-6

* Note that for some electronic prescribing systems three decimal places are not accepted. In these instances the above affected figures can be rounded up (e.g. to 0.73, 1.13, 1.58 and 1.78mg). The resulting doses are not perfectly measurable but will be well within acceptable variance limits.



Drug- and Dose-specific Tables

These tables are intended to assist with non-electronic systems where protocol-specific information is required. The range of the tables has been extended beyond 'normal' use, especially in the lower ranges, to allow for up to a 50% dose reduction to be used in patients who are $1.3m^2$. Doses which are normally capped are highlighted in grey.

Bendamustine

BSA	Choose nearest for dose reductions							1.0	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0
60 mg/m ²	32.5	35	37.5	40	45	50	55	60	67.5	75	75	82.5	90	100	100	112.5	112.5	125	125	125	140	140	157.5	157.5	157.5	175	175	175
70 mg/m ²	35	37.5	40	45	50	55	60	67.5	75	82.5	90	100	100	112.5	125	125	140	140	140	157.5	157.5	175	175	175	197.5	197.5	197.5	222.5
90 mg/m ²	45	50	55	60	67.5	75	82.5	90	100	112.5	112.5	125	140	140	157.5	157.5	175	175	197.5	197.5	197.5	222.5	222.5	222.5	250	250	250	282.5

Bortezomib SC

BSA	Choose nearest for dose reductions							1.0	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0
1.3 mg/m ²	0.65	0.7	0.75	0.85	0.95	1.05	1.15	1.3	1.45	1.6	1.6	1.8	2	2	2.25	2.25	2.5	2.5	2.75	2.75	3	3	3.25	3.5	3.5	3.75	3.75	4

Table usage notes: 'Single container' tables are for products which are able to be supplied as a single unit, such as an infusion bag, one syringe, or as an infusor device (pump). 'Multiple syringe' (e.g. 'pick-and-mix') tables are for products which are supplied in combinations of syringes to make up a single dose. Some drugs may have both a single container table, and a multiple syringe table. Doses may vary between the two tables due to rounding requirements.

