

# **2019/20 National Tariff Payment System: Annex E**

## **Technical guidance for mental health clusters**

**A joint publication by  
NHS England and NHS Improvement**

March 2019

# **2019/20 National Tariff Payment System: Annex E – Technical guidance for mental health clusters**

**A joint publication by NHS England and NHS Improvement**

Version number: 1

First published: March 2019

Updated:

Prepared by: NHS England and NHS Improvement joint pricing team

Classification: OFFICIAL

This information can be made available in alternative formats upon request. Please contact [pricing@improvement.nhs.uk](mailto:pricing@improvement.nhs.uk)

## Contents

1	Introduction .....	5
1.1	What is a cluster? .....	5
1.2	When should I cluster someone? .....	6
1.3	How do I cluster someone who is newly referred? .....	6
1.4	Care review and the clustering process .....	7
1.5	Care transition protocols.....	8
1.6	Step-by-step guide to the use of MHCT ratings, cluster profiles and care transition protocols at care reviews .....	9
1.7	Patient safety.....	10
1.8	Bipolar advice.....	10
1.9	Advice concerning the transition of patients into and out of clusters 14 and 15 (psychotic crisis and psychotic depression) .....	10
	Appendix 1: Mental health clustering tool, version 5.0 .....	12
	Mental health clustering tool (MHCT) version 5.0 (2019/20) .....	12
	Mental health clustering tool (MHCT).....	13
	PART 1: Current ratings.....	14
	PART 2: Historical ratings .....	19
	Appendix 2: Decision tree .....	22
	Appendix 3: Cluster descriptions and care transition protocols .....	23
	Guide to Appendix 3.....	23
	CARE CLUSTER 0: Variance .....	24
	CARE CLUSTER 1: Common mental health problems (low severity).....	25
	CARE CLUSTER 2: Common mental health problems (low severity with greater need) .....	27
	CARE CLUSTER 3: Non-psychotic (moderate severity) .....	29
	CARE CLUSTER 4: Non-psychotic (severe).....	31
	CARE CLUSTER 5: Non-psychotic disorders (very severe) .....	33
	CARE CLUSTER 6: Non-psychotic disorder of over-valued ideas.....	35
	CARE CLUSTER 7: Enduring non-psychotic disorders (high disability).....	37
	CARE CLUSTER 8: Non-psychotic chaotic and challenging disorders.....	39
	CARE CLUSTER 9: Blank cluster .....	41
	CARE CLUSTER 10: First episode psychosis (with/without manic features) ....	43
	CARE CLUSTER 11: Ongoing recurrent psychosis (low symptoms) .....	45
	CARE CLUSTER 12: Ongoing or recurrent psychosis (high disability) .....	47
	CARE CLUSTER 13: Ongoing or recurrent psychosis (high symptom and disability) .....	49
	CARE CLUSTER 14: Psychotic crisis.....	51
	CARE CLUSTER 15: Severe psychotic depression.....	53
	CARE CLUSTER 16: Psychosis and affective disorder (high substance misuse and engagement).....	55
	CARE CLUSTER 17: Psychosis and affective disorder – difficult to engage ....	57

CARE CLUSTER 18: Cognitive impairment (low need) .....	59
CARE CLUSTER 19: Cognitive impairment or dementia complicated (moderate need).....	61
CARE CLUSTER 20: Cognitive impairment or dementia complicated (high need).....	63
CARE CLUSTER 21: Cognitive impairment or dementia (high physical or engagement).....	65
Appendix 4 – Practice guidance note regarding: allocation of patients with a bipolar diagnosis to a cluster.....	67
Summary.....	67
Background.....	67
Bipolar diagnosis.....	68
Current state .....	68
Current generic guidance.....	68
Case vignettes .....	69

# 1 Introduction

The mental health (MH) currencies have been mandated for use since April 2012. For most provider and commissioning organisations completeness and accuracy of cluster allocations is now a key concern and a great deal of audit/assurance work is being undertaken. This booklet is not intended to replace face-to-face training sessions, but to provide clinicians with all the information needed to accurately use the model.

## 1.1 What is a cluster?

In this context, a cluster is a global description of a group of people with similar characteristics as identified from a holistic assessment and then rated using the Mental Health Clustering Tool (MHCT). The clusters allow for a degree of variation in the combination and severity of rated needs. However, as the clusters are statistically underpinned, definite patterns in the MHCT ratings exist for each of them. These ranges are indicated by the colour coded grids ([Appendix 3](#)) and are supplemented by the contextual information on the left-hand side of each page. This is particularly useful when reviewing the appropriateness of previous cluster allocations.

The following list of clusters includes links to the cluster description in Appendix 3

- [CARE CLUSTER 0: Variance](#)
- [CARE CLUSTER 1: Common mental health problems \(low severity\)](#)
- [CARE CLUSTER 2: Common mental health problems \(low severity with greater need\)](#)
- [CARE CLUSTER 3: Non-psychotic \(moderate severity\)](#)
- [CARE CLUSTER 4: Non-psychotic \(severe\)](#)
- [CARE CLUSTER 5: Non-psychotic disorders \(very severe\)](#)
- [CARE CLUSTER 6: Non-psychotic disorder of over-valued ideas](#)
- [CARE CLUSTER 7: Enduring non-psychotic disorders \(high disability\)](#)
- [CARE CLUSTER 8: Non-psychotic chaotic and challenging disorders](#)
- [CARE CLUSTER 9: Blank cluster](#)
- [CARE CLUSTER 10: First episode psychosis \(with/without manic features\)](#)
- [CARE CLUSTER 11: Ongoing recurrent psychosis \(low symptoms\)](#)
- [CARE CLUSTER 12: Ongoing or recurrent psychosis \(high disability\)](#)

- [CARE CLUSTER 13: Ongoing or recurrent psychosis \(high symptom and disability\)](#)
- [CARE CLUSTER 14: Psychotic crisis](#)
- [CARE CLUSTER 15: Severe psychotic depression](#)
- [CARE CLUSTER 16: Psychosis and affective disorder \(high substance misuse and engagement\)](#)
- [CARE CLUSTER 17: Psychosis and affective disorder – difficult to engage](#)
- [CARE CLUSTER 18: Cognitive impairment \(low need\)](#)
- [CARE CLUSTER 19: Cognitive impairment or dementia complicated \(moderate need\)](#)
- [CARE CLUSTER 20: Cognitive impairment or dementia complicated \(high need\)](#)
- [CARE CLUSTER 21: Cognitive impairment or dementia \(high physical or engagement\)](#)

## **1.2 When should I cluster someone?**

People's needs change over time, and over the course of their treatment. A payment system for mental healthcare must reflect the differing levels of input that are provided throughout changing and unpredictable episodes of care. To achieve this, people should not only be assessed and clustered at the point of referral, but also re-assessed and re-clustered periodically. In practice, this will equate to assessing and clustering people at:

- the end of the initial assessment (typically within two contacts)
- all planned Care Programme Approach (CPA)<sup>1</sup> or other formal care reviews
- any other point where a significant change in planned care is deemed necessary (eg unplanned reviews, urgent admissions, etc)

Organisations should ensure there is clarity about who is responsible for clustering, particularly when more than one professional is involved.

## **1.3 How do I cluster someone who is newly referred?**

As organisations use different IT systems, the exact procedures will vary from provider to provider. However, all providers will follow these basic steps:

---

<sup>1</sup> CPA provides people with enduring mental health conditions with a care co-ordinator and a care plan which must be regularly reviewed.

- **Step 1:** Based on the information you have gathered during your routine screening/assessment process, rate the individual's identified needs using the Mental Health Clustering Tool - Version 5.0 ([Appendix 1](#)).
- **Step 2:** Use the decision tree ([Appendix 2](#)) to decide if the presenting needs are non-psychotic, psychotic or organic in origin. Then decide which of the next level of headings is most accurate. This will have narrowed down the list of clusters that are likely to describe the person's needs.
- **Step 3:** Look at the rating grids ([Appendix 3](#)) to decide which one is the most appropriate by using the colour-coded key.
  - Start with the red ratings. These indicate the type and level of need which must be apparent to be a member of this cluster. If the ratings do not match, try another cluster.
  - Next, consider the orange ratings. These represent expected ratings. You may allocate a person to a cluster if the orange ratings do not exactly match the coloured grids. However, this reflects a 'weaker fit' to that cluster.
  - Finally review the yellow ratings. These represent ratings that may occur. These scales have significantly less bearing on cluster allocation but may indicate the need for additional care plan interventions.

**Remember, the final clustering decision is yours, based on your assessment results and your clinical judgement in applying this guidance.**

#### **1.4 Care review and the clustering process**

Every day, practitioners make decisions about starting, stopping, increasing and decreasing interventions. These decisions are made according to a range of complex and inter-related factors, but primarily in response to individual service user need. The currency model describes these individually assessed needs in a consistent way, through use of the MHCT and the resulting set of needs-based clusters.

The clusters, therefore, describe groups of service users with similar types of characteristics. These groups/clusters can be compared to each other in a variety of ways including: severity of need; complexity of need; acuity; intensity of likely treatment response; anticipated course of illness, etc.

While some comparisons will be more useful than others in different situations, in this booklet a global judgement is made which combines all these factors and leads

either to the term ‘**step-up**’ or the term ‘**step-down**’ being used to describe movement between any given clusters.

### 1.5 Care transition protocols

The points at which the appropriateness of the current cluster allocation is reconsidered should not be arbitrary. It should occur at natural and appropriate points in the individual’s care pathway. Typically, these are termed reviews but it is important to note that reviews can be relatively informal as well as formal, and can be in response to unforeseen changes in need – ie, unplanned as well as pre-planned.

Consider the following clinical scenarios:

- The planned review of a service user halfway through a course of 16 sessions of cognitive behaviour therapy (CBT) for depression will often reveal significant improvements and a corresponding reduction in MHCT ratings for anxiety and low mood. This is rarely seen as a sustainable change in the user’s presentation and thus the original treatment plan continues until the intervention is completed, rather than be reduced to a lower intensity intervention (eg computerised CBT).
- Some months after treatment from an Assertive Outreach Team begins, improvements in presentation (particularly patterns of engagement) are not uncommon. These are unlikely to trigger a significant reduction in the overall level of intervention provided until the improvements have been maintained for some time. Therefore, the cluster allocation that originally triggered an assertive and intensive service response remains valid, as it is still seen as a truer reflection of the individual’s overall needs.
- Service users diagnosed with borderline personality disorder are well known to exhibit erratic patterns of behaviour, with fluctuations in distress and risk commonplace. Despite increases in risk, decisions are often made to take therapeutic risks rather than immediately increasing the overall level of intervention in response to what may turn out to be transient and self-limiting increases in perceived need.

From these examples it is clear individuals only fit the needs profiles for the appropriate cluster at certain key points in their journey (ie the start of a period of care). At clinical reviews, additional factors must also be taken into account before an alternative cluster allocation is made and care is changed significantly.



These factors are described in this booklet as care transition protocols and include the step-up and step-down criteria for each cluster. **Only when a set of criteria has been met should the allocated cluster be changed to that suggested by the clustering tool ratings.** The protocols also include examples of local discharge criteria which outline the circumstances when service users could be discharged from in-scope mental health services completely. **Please note:** providers and commissioners will need to agree their own local discharge criteria.

The care transition pages in this booklet describe, for each cluster:

- the length of time service users are likely to remain in mental health (MH) services
- a frequency for re-assessing the appropriateness of the cluster
- the likelihood of each possible cluster transition.

It also attempts to visually represent the relationship between each cluster in terms of intensity, acuity, etc. **Please note:** in general, cluster reviews should be aligned to care reviews. The review frequencies quoted are outer limits, not absolute frequencies.

As most practitioners work with specific groups of service users and will only encounter a small number of clusters routinely, they should become familiar with the sections of this booklet that are relevant to their work. In addition, the six steps described below guide practitioners through the process.

## **1.6 Step-by-step guide to the use of MHCT ratings, cluster profiles and care transition protocols at care reviews**

- 1 Select the page containing care transition protocols that correspond to the individual's current cluster.
- 2 After completing an appropriate re-assessment of risks and needs, complete a new MHCT.
- 3 Consider the **step-up criteria**. If any one of these is met, this suggests the current cluster allocation needs to change and, with reference to the clustering booklet, the latest MHCT ratings should be used to decide on the new cluster. If the step-up criteria are not met...
- 4 Consider the **discharge criteria**. If all of these are met, this indicates the need to explore discharge from in-scope mental health services back to GP-led (primary) care. If the discharge criteria are not met...

- 5 Consider the **step-down criteria**. If all of these are met, this suggests the current cluster allocation needs to change and, with reference to the clustering booklet, the MHCT ratings should be used to decide on the new cluster. If the step-down criteria are not met...
- 6 This indicates that the existing cluster allocation remains valid, as any differences in the user's needs that have occurred do not warrant the changes in service response that allocation to a different cluster would trigger.

## 1.7 Patient safety

Any issues relating to service user safety that arise through the use of the MHCT and the mental health care clusters should be raised through your organisation's own patient safety reporting routes. Any urgent service user safety issues that directly relate to the clustering tool or the clusters should also be reported via [pricing@improvement.nhs.uk](mailto:pricing@improvement.nhs.uk)

## 1.8 Bipolar advice

All patients, including those with a diagnosis of bipolar disorder, should be allocated to the cluster **which best describes the combination and severity of their primary presenting needs**. Patients with the same diagnosis can therefore be accurately allocated to different clusters within a superclass (non-psychotic, psychotic or organic). As a diagnosis of bipolar disorder covers a particularly wide variety of presentations, these patients may be allocated to either a psychotic or a non-psychotic cluster depending on your clinical judgement. However, consideration of the likely and unlikely diagnoses sections of the clustering booklet should be carefully considered.

Please see [Appendix 4](#) for further practice guidance regarding allocating patients with bipolar diagnosis to a cluster.

## 1.9 Advice concerning the **transition of patients into and out of clusters 14 and 15 (psychotic crisis and psychotic depression)**

Patients' needs change over time. The clusters are relatively broad in nature and hence there will be a range of presentations (more to less well) within each. It is not unusual for patients in clusters 10, 13, 16 and 17 to relapse and be better described by cluster 14 (ie be experiencing a psychotic crisis). In this case, assuming the change in presentation warrants a significant change in the intensity of their treatment package, they should be re-clustered accordingly (ie to cluster 14). While not universally followed, this process is reasonably well recognised. On recovery,

however, it is important to **take account of the patient's previous presentation** when stepping the patient down from cluster 14 post-crisis. So if, for example, the patient has been in cluster 17 before their psychotic crisis, even if their symptoms are well controlled, at the point of stepping down from 14 it is unlikely that anything other than cluster 17 will meet their needs. Similarly, if a patient in cluster 10 (first episode psychosis) becomes acutely unwell and warrants allocation to cluster 14, on stepping-down they should be re-allocated to cluster 10 for the rest of their three-year treatment package if they are within three years of initial presentation.

## Appendix 1: Mental health clustering tool, version 5.0

### Mental health clustering tool (MHCT) version 5.0 (2019/20)

The MHCT incorporates items from the Health of the Nations Outcome Scales (HoNOS), (Wing et al 1999<sup>2</sup>) and the Summary of Assessments of Risk and Need (SARN), (Self et al 2008<sup>2</sup>) to provide all the information necessary to allocate individuals to clusters.

#### HoNOS

HoNOS is an internationally recognised outcome measure developed by the Royal College of Psychiatrists Research Unit (CRU) to measure health and social functioning outcomes in Mental Health Services. The aim of HoNOS was to produce a brief measure capable of being completed routinely by clinicians and recorded as part of a minimum mental health dataset. The first 12 items of the MHCT are HoNOS items. The HoNOS items are used here with the permission of the Royal College of Psychiatrists, which holds the copyright. A review of HoNOS was concluded in 2018, and the results adopted by the Royal College of Psychiatrists and their international partners.<sup>3</sup> This has resulted in changes to the HoNOS glossary and these changes are detailed in the Current Ratings table for items 1-12.

#### SARN

The Summary of Assessments of Risk and Need (SARN) was developed by the Care Pathways and Packages Project<sup>45</sup> to aid the process of establishing a classification of service users based on their needs so that appropriate service responses could be developed at both individual and service level. It provides a brief description of the needs of people entering mental health services for the first time or presenting with a possible need for change in their care or treatment. It allows professionals from a range of backgrounds to summarise their assessments in a shared format. Thus it provides a common language for describing health states and related social conditions and improves communication between different users of the tool including health and social care professionals, Service Users themselves, commissioners and researchers.

---

<sup>2</sup> Wing, J. K., Curtis, R. H. & Beevor, A. S. (1999) Health of the Nation Outcome Scales (HoNOS). *British Journal of Psychiatry*, 174 (5), 432-434.

<sup>3</sup> [www.cambridge.org/core/journals/bjpsych-bulletin/article/review-and-update-of-the-health-of-the-nation-outcome-scales-honos/85ED388E4268748F59BCB4BCC43C7BEE/core-reader](http://www.cambridge.org/core/journals/bjpsych-bulletin/article/review-and-update-of-the-health-of-the-nation-outcome-scales-honos/85ED388E4268748F59BCB4BCC43C7BEE/core-reader)

<sup>4</sup> Self R; Rigby A; Leggett C and Paxton R (2008) Clinical Decision Support Tool: A rational needs-based approach to making clinical decisions. *Journal of Mental Health*, 17(1): 33-48.

<sup>5</sup> CPPP was a consortium of providers and commissioners in Yorkshire and the North East that undertook the initial development work on mental health currencies

## Mental health clustering tool (MHCT)

**Part 1** contains scales relating to the severity of problems experienced by the individual during the two weeks before the date of the rating.

**Part 2** contains scales that consider problems from a 'historical' perspective. These will be problems that occur in episodic or unpredictable ways. While they may not have been experienced by the individual during the two weeks before the rating date, clinical judgement would suggest that there is still a cause for concern that cannot be disregarded (ie no evidence to suggest that the person has changed since the last occurrence as a result of time, therapy, medication or environment, etc). In these circumstances, any event that remains relevant to the cluster allocation (and hence the interventions offered) should be included.

### *Summary of rating information*

- Rate each scale in order from 1 to 13 (Part 1), followed by A to E (Part 2).
- For the first 12 scales, do not include information rated in an earlier scale except for scale 10 which is an overall rating.
- Rate the MOST SEVERE problem that occurred in the rating period
- All scales follow the format:
  - 0 = no problem
  - 1 = minor problem requiring no action
  - 2 = mild problem but definitely present
  - 3 = moderately severe problem
  - 4 = severe to very severe problem
- Record 9 if Not Known but be aware that this is likely to make accurate clustering impractical and indicate that further assessment is required.

## PART 1: Current ratings

For scales 1-13, rate the most severe occurrence in the previous two weeks

1. Overactive, aggressive, disruptive or agitated behaviour (current)					
	0	1	2	3	4
<ul style="list-style-type: none"> <li>Rate any of the behavioural components that this scale covers from overactive or aggressive or disruptive or agitated behaviours.</li> <li>Include such behaviour due to any cause (e.g. drugs; alcohol; dementia; psychosis; depression).</li> <li>Do not include bizarre behaviour to be rated at Scale 6, unless it is aggressive, destructive or overactive.</li> </ul>	No problem of this kind during the period rated.	Irritability, quarrels, restlessness, etc, not requiring action.	Includes aggressive gestures, pushing or pestering others; threats or verbal aggression; lesser damage to property (eg broken cup, window); marked over-activity or agitation.	Physically aggressive to others or animals (short of rating 4); threatening manner; more serious over-activity or destruction of property.	At least one serious physical attack on others or on animals; destructive of property (eg fire-setting); serious intimidation or obscene behaviour.
2. Non-accidental self-injury (current)					
	0	1	2	3	4
<ul style="list-style-type: none"> <li>Do not include accidental self-injury (due e.g. to dementia or severe learning disability); the cognitive problem is to be rated at Scale 4 and the injury at Scale 5.</li> <li>Do not include illness or injury as a direct consequence of drug/alcohol use (e.g. cirrhosis of the liver or injury resulting from drink driving) to be rated at Scale 5.</li> </ul>	No problem of this kind during the period rated.	Fleeting thoughts about ending it all but little risk during the period rated; no self-harm.	Mild risk during the period rated; includes more frequent thoughts or talking about self-harm or suicide (including 'passive' ideas of self-harm such as not taking avoiding action in a potentially life threatening situation e.g. while crossing a road).	Moderate to serious risk of deliberate self-harm; includes frequent/persistent thoughts or talking about self-harm; includes preparatory behaviours (e.g. collecting tablets)	Serious suicidal attempt and/or serious deliberate self-injury during the period rated.
3. Problem-drinking or drug-taking (current)					
	0	1	2	3	4
<ul style="list-style-type: none"> <li>Include psychological as well as behavioural impacts of drug (illicit and/or prescription) and alcohol use.</li> <li>Do not include aggressive/destructive behaviour due to alcohol or drug use already rated at Scale 1.</li> <li>Do not include physical illness or disability due to alcohol or drug use to be rated at Scale 5.</li> <li>Do not include dependence on tobacco products unless there are severe and adverse consequences arising from that addiction above and beyond the known long-term harms to physical health.</li> </ul>	No problem of this kind during the period rated.	Some excessive consumption but no adverse consequences	Excessive and/or harmful consumption resulting in adverse consequences, but no obvious craving or dependency.	Definite craving and/or dependence on alcohol or drugs.	Severe craving/dependence resulting in severe adverse consequences from alcohol/drug problems.

4. Cognitive problems (current)					
	0	1	2	3	4
<ul style="list-style-type: none"> <li>Include problems of orientation, memory, language, thought disorder and problem solving associated with any disorder: dementia, learning disability, schizophrenia, etc.</li> <li>Do not include temporary problems (e.g. hangovers) which are clearly associated with alcohol, drug or medication use, rated at Scale 3.</li> <li>Do not rate disorders of thought content (e.g. eccentric or delusional thinking) that will be rated at Scale 6.</li> <li></li> </ul>	No problem of this kind during the period rated.	Minor problems with orientation (e.g. occasionally disorientated to time); memory (e.g. occasionally forgets names); language (e.g. on occasions unable to clearly express ideas; or has to have questions and instructions repeated); problem solving (e.g. able to solve simple problems but some difficulty with complex tasks).	Mild but definite problems with orientation (e.g. lost way in an unfamiliar place); memory (e.g. some difficulty remembering events; learning new material); language (e.g. some difficulty understanding and/or expressing ideas); mild thought disorder; problem solving (e.g. sometimes mixed up about simple decisions.)	Moderate problems with orientation (e.g. lost way in a familiar place; often disorientated to time); memory (e.g. new material rapidly lost; only highly learned material retained); language (e.g. speech can be incoherent; fails to understand common words/phrases); moderate thought disorder evident; problem solving (e.g. frequently unable to think clearly or solve simple problems).	Severe difficulties with orientation (e.g. consistently disorientated to time, person and place); memory (e.g. loss of distant and recent memory; unable to learn new information); language (e.g. very limited receptive or expressive communication); severe thought disorder; no effective problem solving.
5. Physical illness or disability problems (current)					
	0	1	2	3	4
<ul style="list-style-type: none"> <li>Include illness or disability from any cause that limits or prevents movement, or impairs sight or hearing, or otherwise interferes with personal functioning.</li> <li>Include side-effects from medication; effects of drug/alcohol use; physical disabilities resulting from accidents or self-harm associated with cognitive problems, drink-driving, etc.</li> <li>Do not include mental/behavioural problems rated at Scale 4.</li> </ul>	No physical health problem during the period rated.	Minor health problems during the period (eg cold, non-serious fall, etc.)	Physical health problem imposes mild restriction on mobility and activity.	Moderate degree of restriction on activity due to physical health problem.	Severe or complete incapacity due to physical health problem.
6. Problems associated with hallucinations and delusions (current)					
	0	1	2	3	4
<ul style="list-style-type: none"> <li>Include hallucinations and/or delusions irrespective of diagnosis.</li> <li>Include unusual and bizarre behaviour associated with hallucinations or delusions.</li> <li>Do not include aggressive, destructive or overactive behaviours attributed to hallucinations and/or delusions, already rated at Scale 1.</li> </ul>	No evidence of hallucinations or delusions during the period rated.	Somewhat unusual or eccentric beliefs not in keeping with cultural norms.	Hallucinations or delusions are present, but there is little distress to patient or manifestation in bizarre behaviours, i.e. clinically present but mild.	Marked preoccupation with hallucinations or delusions, causing much distress and/or manifested in obviously bizarre behaviour, i.e. moderately severe clinical problem.	Mental state and behaviour is seriously and adversely affected by hallucinations or delusions, with severe impact on patient.
7. Problems with depressed mood (current)					
	0	1	2	3	4
<ul style="list-style-type: none"> <li>Include cognitive, affective or behavioural aspects of depressed mood (e.g. loss of interest or pleasure; lack of energy; loss of self-esteem; feelings of guilt).</li> </ul>	No problem associated with depressed mood during the period rated.	Gloomy or minor changes in mood.	Mild but definite depressed mood and	Moderate depressed mood on subjective or objective measures	Severe depressed mood on subjective or objective grounds (e.g.

<ul style="list-style-type: none"><li>Do not include overactivity or agitation, already rated at Scale 1.</li><li>Do not include suicidal ideation or attempts, already rated at Scale 2.</li><li>Do not include delusions or hallucinations, already rated at Scale 6.</li><li>Do not include other symptoms of depression as described at Scale 8 (i.e. changes in sleep, appetite or weight; anxiety symptoms).</li><li></li></ul>			distress (e.g. loss of interest or pleasure; feelings of guilt; loss of self-esteem).	(depressive symptoms more marked).	profound loss of interest or pleasure; preoccupation with ideas of guilt or worthlessness).										
8. Other mental and behavioural problems (current)															
	0	1	2	3	4										
<ul style="list-style-type: none"><li>Rate only the most severe mental and behavioural problem not considered in previous items.</li><li>Specify the type of problem by entering the appropriate letter from the following table.</li></ul> <table><tr><td>A Phobic: Fear or avoidance behaviour in response to specific situations/objects that is out of proportion to actual threat.</td></tr><tr><td>B Anxiety: Patient experiences general anxiety, panic or similar experiences.</td></tr><tr><td>C Obsessive-compulsive: Recurrent obsessions or compulsive acts that are distressing and typically perceived by the patient as irrational.</td></tr><tr><td>D Reactions to stressful events or trauma: Acute stress reactions and/or response to traumatic events.</td></tr><tr><td>E Dissociative: Mental process where the patient disconnects from their thoughts, feelings, memories or sense of identity.</td></tr><tr><td>F Somatoform: Persistent perceived physical health problems that have no known medical basis.</td></tr><tr><td>G Eating: Excessive intake or persistent restriction of food intake; includes related disordered behaviours to manage weight e.g. purging, excessive exercise, dieting etc.</td></tr><tr><td>H Sleep: Problems with the quality, timing or duration of sleep that impact on sense of fatigue, cognitive function or mood.</td></tr><tr><td>I Sexual: Disturbance of the patient's ability to respond sexually or experience sexual pleasure.</td></tr><tr><td>K Elated mood: Feelings of euphoria, excitement, expansive mood or optimism that do not reflect person's actual circumstances.</td></tr></table>	A Phobic: Fear or avoidance behaviour in response to specific situations/objects that is out of proportion to actual threat.	B Anxiety: Patient experiences general anxiety, panic or similar experiences.	C Obsessive-compulsive: Recurrent obsessions or compulsive acts that are distressing and typically perceived by the patient as irrational.	D Reactions to stressful events or trauma: Acute stress reactions and/or response to traumatic events.	E Dissociative: Mental process where the patient disconnects from their thoughts, feelings, memories or sense of identity.	F Somatoform: Persistent perceived physical health problems that have no known medical basis.	G Eating: Excessive intake or persistent restriction of food intake; includes related disordered behaviours to manage weight e.g. purging, excessive exercise, dieting etc.	H Sleep: Problems with the quality, timing or duration of sleep that impact on sense of fatigue, cognitive function or mood.	I Sexual: Disturbance of the patient's ability to respond sexually or experience sexual pleasure.	K Elated mood: Feelings of euphoria, excitement, expansive mood or optimism that do not reflect person's actual circumstances.	No evidence of any of these problems during period rated.	Minor non-clinical problems only.	A problem is clinically present, but at a mild level (e.g. the problem is intermittent; the patient maintains a degree of control or is not unduly distressed).	Moderately severe clinical problem (e.g. more frequent, more distressing or more marked symptoms).	Severe problem which dominates or seriously affects many activities.
A Phobic: Fear or avoidance behaviour in response to specific situations/objects that is out of proportion to actual threat.															
B Anxiety: Patient experiences general anxiety, panic or similar experiences.															
C Obsessive-compulsive: Recurrent obsessions or compulsive acts that are distressing and typically perceived by the patient as irrational.															
D Reactions to stressful events or trauma: Acute stress reactions and/or response to traumatic events.															
E Dissociative: Mental process where the patient disconnects from their thoughts, feelings, memories or sense of identity.															
F Somatoform: Persistent perceived physical health problems that have no known medical basis.															
G Eating: Excessive intake or persistent restriction of food intake; includes related disordered behaviours to manage weight e.g. purging, excessive exercise, dieting etc.															
H Sleep: Problems with the quality, timing or duration of sleep that impact on sense of fatigue, cognitive function or mood.															
I Sexual: Disturbance of the patient's ability to respond sexually or experience sexual pleasure.															
K Elated mood: Feelings of euphoria, excitement, expansive mood or optimism that do not reflect person's actual circumstances.															



<p>O Other: Any other mental or behavioural problem, not rated elsewhere, that is significant that results in patient distress or impacts upon their behaviour.</p>					
<p>N.B. J has been deliberately omitted to allow compatibility with the previous version of the HoNOS.</p>					
<b>9. Problems with relationships (current)</b>					
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
<ul style="list-style-type: none"> <li>Rate the patient's most severe problem associated with active or passive withdrawal from interpersonal relationships, and/or non-supportive, destructive or self-damaging relationships.</li> <li>Include family as well as broader social relationships.</li> </ul>	No significant problem during the period.	Minor non-clinical problems.	Definite problems in making or sustaining supportive relationships which create mild but significant distress or difficulty for the patient; patient complains and/or problems are evident to others.	Persisting major problems due to active or passive withdrawal from social relationships; relationships that provide little or no comfort or support; and/or problematic relationships which create moderate levels of distress or difficulty for the patient.	Severe and distressing social isolation or withdrawal from social relationships; and/or problematic relationships which create severe levels of distress or difficulty for the patient.
<b>10. Problems with activities of daily living (current)</b>					
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
<ul style="list-style-type: none"> <li>Rate the overall level of functioning in activities of daily living (ADL) (e.g. problems with basic activities or self-care such as eating, washing, dressing, toilet; also complex skills such as budgeting, organising where to live, occupation and recreation, mobility and use of transport, shopping, self-development, etc.).</li> <li>Rate what the person is capable of doing, independently of current support from others.</li> <li>Include any lack of motivation, including the use of self-help opportunities, since this contributes to a lower overall level of functioning.</li> <li>Do not include lack of opportunities for exercising intact abilities and skills, to be rated at Scales 11-12.</li> </ul>	No problem during period rated; good ability to function in all areas.	Minor problems only with self-care without significantly adverse consequences (e.g. untidy; disorganised), and / or minor difficulty with complex skills but still able to function independently.	Self-care and basic activities adequate (though some prompting may be required) but major lack of performance of one or more complex skills (see above).	Major problems in one or more areas of self-care (e.g. eating; washing; dressing; toilet) as well as major inability to perform several complex skills.	Severe disability or incapacity in all or nearly all areas of self-care and complex skills.
<b>11. Problems with living conditions (current)</b>					
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
<p><b>NB: Rate patient's usual housing and living conditions.</b> In general, try to rate the housing and living conditions most relevant to the patient's situation (e.g. if a brief stay in an acute ward is anticipated, rate the patient's home environment; if discharge is imminent, rate the patient's destination accommodation; if a lengthy hospital stay (e.g. over 6 months) is anticipated, rate the suitability of the ward).</p> <ul style="list-style-type: none"> <li>Rate the overall severity of problems with the quality of housing and living conditions. Are the basic necessities</li> </ul>	Housing and living conditions are acceptable; helpful in keeping any disabilities rated at Scale 10 to the lowest level possible and supportive of self-help.	Housing and living conditions are reasonably acceptable although there are minor or transient problems (e.g. not ideal location; not preferred option etc.).	Problem with one or more aspects of housing or living conditions (e.g. limited facilities to improve patient's independence).	Multiple significant problems with housing or living conditions (e.g. some basic necessities absent; housing or living conditions have minimal or no facilities to improve patient's independence).	Housing or living conditions are unacceptable (e.g. lack of basic necessities; patient is at risk of eviction or 'roofless'; or living conditions are otherwise intolerable) making patient's problems worse.

<p>met (e.g. adequate heat; light; sanitation; cooking facilities)?</p> <ul style="list-style-type: none"> <li>In addition to basic necessities, consider how well the patient's housing and living conditions match their current needs.</li> </ul> <p>Do not rate the level of functional disability itself, already rated at Scale 10.</p> <p>If information not available, rate at Scale 9.</p>					
<b>12. Problems with occupation and activities (current)</b>					
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
<ul style="list-style-type: none"> <li><b>NB: Rate patient's usual situation.</b> In general, try to rate the occupation and activities most relevant to the patient's situation (e.g. if a brief stay in an acute ward is anticipated, rate the patient's usual occupation and activities; if discharge is imminent, rate the occupation and activities of the patient's destination; if a lengthy hospital stay (e.g. over 6 months) is anticipated, rate the suitability of the ward's provision).</li> <li>Rate the overall level of problems with the quality of meaningful occupation and activities. Is there help to cope with disabilities, and opportunities for maintaining or improving occupational and recreational skills and activities? Consider factors such as stigma; lack of suitably skilled staff; access to supportive facilities (e.g. staffing and equipment of day centres, workshops, social clubs, etc.).</li> <li>Consider how well the patient's occupation and activities match their current needs.</li> </ul> <p>Do not rate the level of functional disability itself, already rated at Scale 10.</p>	<p>Patient's occupation and activities are acceptable; helpful in keeping any disability rated at Scale 10 to the lowest level possible and supportive of self-help, and maximising autonomy and role functioning.</p>	<p>Minor or temporary problems (e.g. reasonable facilities available but not always at desired times, etc.).</p>	<p>Limited choice of activities to maintain or improve autonomy and role functioning (e.g. there is a lack of reasonable tolerance such as unfairly refused entry to public facilities; or insufficient skilled services; or helpful service is available but for very limited hours).</p>	<p>Marked deficiency in skilled services available to help minimise level of disability and help optimise autonomy and role functioning. No opportunities to use intact skills or add new ones; unskilled care difficult to access.</p>	<p>Lack of any opportunity for meaningful activities, or complete inability of services to involve the patient in such activities, may make patient's problems worse.</p>
<b>13. Strong unreasonable beliefs that are <u>not</u> psychotic in origin (current)</b>					
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
<ul style="list-style-type: none"> <li>Rate any apparent strong unreasonable beliefs (found in <b>some</b> people with disorders such as obsessive compulsive disorder, anorexia nervosa, personality disorder, morbid jealousy, etc)</li> <li>Do not include delusions rated at scale 6.</li> <li>Do not include severity of disorders listed above where strong unreasonable beliefs are not present – rated at Scale 8.</li> <li>Do not include beliefs/behaviours consistent with a person's culture.</li> </ul>	<p>No strong unreasonable beliefs evident.</p>	<p>Holds illogical or unreasonable belief(s) but has insight into their lack of logic or reasonableness and can challenge them most of the time and they have only a minor impact on the individual's life.</p>	<p>Holds illogical or unreasonable belief(s) but individual has insight into their lack of logic or reasonableness. Belief(s) can be successfully challenged by individual on occasions. Beliefs have a mild impact on the person's life.</p>	<p>Holds strong illogical and unreasonable belief(s) but has some insight into the relationship between the beliefs and the disorder. Belief(s) can be 'shaken' by rational argument. Tries to resist belief but with little effect. Has a significant negative impact on person's life. he disorder makes treatment more difficult than usual.</p>	<p>Holds strong illogical or unreasonable belief(s) with little or no insight in the relationship between the belief and the disorder. Belief(s) cannot be 'shaken' by rational argument. Does not attempt to resist belief(s). Has a significant negative impact on the person's life or other people's lives and the disorder is very resistant to treatment.</p>

## PART 2: Historical ratings

Scales A-E rate problems that occur in an episodic or unpredictable way. Include any event that remains relevant to the current plan of care.

While there may or may not be any direct observation or report of a manifestation during the last two weeks, the evidence and clinical judgement would suggest that there is still a cause for concern that cannot be disregarded (ie no evidence to suggest that the person has changed since the last occurrence as a result of time, therapy, medication or environment, etc).

<b>A. Agitated behaviour/expansive mood (historical)</b>					
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
<ul style="list-style-type: none"> <li>Rate agitation and overactive behaviour causing disruption to social role functioning. Behaviour causing concern or harm to others.</li> <li>Elevated mood that is out of proportion to circumstances.</li> <li>Include such behaviour due to any cause (e.g. drugs, alcohol, dementia, psychosis, depression etc.)</li> <li>Excessive irritability, restlessness, intimidation, obscene behaviour and aggression to people animals or property.</li> <li>Do not include odd or bizarre behaviour to be rated at Scale 6.</li> </ul>	No needs in this area.	Presents as irritable, argumentative with some agitation. Some signs of elevated mood or agitation not causing disruption to functioning.	Makes verbal/gestural threats. Pushes/pesters but no evidence of intent to cause serious harm. Causes minor damage to property (eg glass or crockery). Is obviously over-active or agitated.	Agitation or threatening manner causing fear in others. Physical aggression to people or animals. Property destruction. Serious levels of elevated mood, agitation, restlessness causing significant disruption to functioning.	Serious physical harm caused to persons/animals. Major destruction of property. Seriously intimidating others or exhibiting highly obscene behaviour. Elevated mood, agitation, restlessness causing complete disruption.
<b>B. Repeat self-harm (historical)</b>					
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
<ul style="list-style-type: none"> <li>Rate repeat acts of self-harm with the intention of managing people, stressful situations, emotions or to produce mutilation for any reason.</li> <li>Include self-cutting, biting, striking, burning, breaking bones or taking poisonous substances, etc.</li> <li>Do not include accidental self-injury (due eg to learning disability or cognitive impairment); the cognitive problem is rated at Scale 4 and the injury at Scale 5.</li> <li>Do not include harm as a direct consequence of drug/alcohol use (eg liver damage) to be rated at Scale 3. Injury sustained while intoxicated to be rated at Scale 5.</li> <li>Do not include harm with intention of killing self (rated at Scale 2).</li> </ul>	No problem of this kind.	Superficial scratching or non-hazardous doses of drugs.	Superficial cutting, biting, bruising, etc or small ingestions of hazardous substances unlikely to lead to significant harm even if hospital treatment not sought.	Repeat self-injury requiring hospital treatment. Possible dangers if hospital treatment not sought. However, unlikely to leave lasting severe damage even if behaviour continues providing hospital treatment sought.	Repeat serious self-injury requiring hospital treatment and likely to leave lasting severe damage if behaviour continues (ie severe scarring, crippling or damage to internal organ) and possibly to death.


<b>C. Safeguarding other children and vulnerable adults (historical)</b>					
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
<ul style="list-style-type: none"> <li>Rate the potential or actual impact of the patient's mental illness, or behaviour, on the safety and well-being of vulnerable people of any age.</li> <li>Include any patient who has substantial access and contact with children or other vulnerable persons.</li> <li>Do not include risk to wider population covered at Scale A.</li> <li>Do not include challenge to relationships covered in Scale 9.</li> </ul>	No obvious impact of the individual's illness or behaviour on the safety or well-being of vulnerable persons.	Mild concerns about the impact of the individual's illness or behaviour on the safety or well-being of vulnerable persons.	Illness or behaviour has an impact on the safety or well-being of vulnerable persons. The individual is aware of the potential impact but is supported and is able to make adequate arrangements.	Illness or behaviour has an impact on the safety or well-being of vulnerable persons but does not meet the criteria to rate 4. There may be delusions, non-accidental self-injury risk or self-harm. However, the individual has insight, can take action to significantly reduce the impact of their behaviour on the children and is adequately supported.	Without action the illness or behaviour is likely to have direct or indirect significant impact on the safety or well-being of vulnerable persons. Problems such as delusions, severe non-accidental self-injury risk or problems of impulse control may be present. There may be lack of insight, an inability or unwillingness to take precautions to protect vulnerable persons and/or lack of adequate support and protection for vulnerable persons.
<b>D. Engagement (historical)</b>					
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
<ul style="list-style-type: none"> <li>Rate the individual's motivation and understanding of their problems, acceptance of their care/treatment and ability to relate to care staff.</li> <li>Include the ability, willingness or motivation to engage in their care/ treatment appropriately, agreeing personal goals, attending appointments. Dependency issues.</li> <li>Do not include cognitive issues as in Scale 4, severity of illness or failure to comply due to practical reasons.</li> </ul>	Has ability to engage/disengage appropriately with services. Has good understanding of problems and care plan.	Some reluctance to engage or slight risk of dependency. Has understanding of own problems.	Occasional difficulties in engagement, ie missed appointments or contacting services between appointments inappropriately. Some understanding of own problems.	Contacts services inappropriately. Has little understanding of own problems. Unreliable attendance at appointments. Or attendance depends on prompting or support.	Contacts multiple agencies, ie GP, A&E, etc, constantly. Little or no understanding of own problems. Fails to comply with planned care. Rarely attends appointments. Refuses service input. Or attendance and compliance depends on intensive prompting and support.

<b>E. Vulnerability (historical)</b>					
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
<ul style="list-style-type: none"> <li>Rate failure of an individual to protect themselves from risk of harm to their health and safety or well-being.</li> <li>Include physical, sexual, emotional and financial exploitation or harm/harassment</li> <li>Do not include problems of engagement rated at Scale D.</li> </ul>	No vulnerability evident.	No significant impact on person's health, safety or well-being.	Concern about the individual's ability to protect their health, safety or well-being requiring support or removal of existing support would increase concern.	Clear evidence of significant vulnerability affecting the individual's ability to protect their health and safety or well-being that requires support (but not as severe as a rating of 4). Or removal of existing support would increase risk.	Severe vulnerability – total breakdown in individual's ability to protect themselves resulting in major risk to the individual's health, safety or well-being.

Item	Rating	Item	Rating
Part 1 (Current)		Part 2 (Historical)	
1		A	
2		B	
3		C	
4		D	
5		E	
6			
7			
8			
Please circle	A B C D E F G H I J		
NB: If J – (other) please specify			
9			
10			
11			
12			
13			

## Appendix 2: Decision tree


### DECISION TREE (RELATIONSHIP OF CARE CLUSTERS TO EACH OTHER)





## Appendix 3: Cluster descriptions and care transition protocols

### Guide to Appendix 3



## CARE CLUSTER 0: Variance

Cluster reviews at least every: 6 months

### Description:

Despite careful consideration of all the other clusters, this group of service users is not adequately described by any of their rating profiles or descriptions. They do however require mental health care and will be offered a service.

### Likely primary diagnosis:

### Unlikely primary diagnosis:

### Impairment:

### Risk:

### Course:

### Likely NICE guidance:

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive Problems					
5	Physical Illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children and vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

**Must score**

**Expected to score**

**May score**

**Unlikely to score**

**No data available**

\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.



## CARE CLUSTER 1: Common mental health problems (low severity)

### Description:

This group has definite but minor problems of depressed mood, anxiety or other disorder but they do not present with any distressing psychotic symptoms.

### Likely primary diagnosis:

May not attract a formal diagnosis but may include mild symptoms of: F32 Depressive episode, F40 Phobic anxiety disorders, F41 Other anxiety disorders, F42 Obsessive-compulsive disorder, F43 Stress reaction/adjustment disorder, F50 Eating disorder.

### Unlikely primary diagnosis:

F00-03 Dementias, F20-29 Schizophrenia, schizotypal and delusional disorders, F30 Manic episode, F31 Bipolar disorder, F33 Major depressive disorder, recurrent.

### Impairment:

Disorder unlikely to cause disruption to wider functioning.

### Risk:

Unlikely to be an issue.

### Course:

The problem is likely to be short term and related to life events.

### Likely NICE guidance:

Service user experience in adult mental health CG136, Anxiety CG113, Depression in adults CG90, Depression with Chronic Health Problems CG91, Common mental health disorders CG123, OCD CG31, Eating Disorders CG9.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical Illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children and vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score



Unlikely to score



Expected to score



No data available



May score




\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 1: Common mental health problems (low severity)

Indicative episode of care: 8 – 12 weeks

Cluster reviews at least every: 12 weeks

Cluster	Step-up criteria (The following criterion is met)	<u>Example local discharge criteria for MH services</u> ( <u>All</u> the following criteria are met)	Step-down criteria (The following criterion is met)
1	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>MHCT V1 item 2 (Non-accidental self-injury) = 0</li> <li>MHCT V1 item 7 (Depression) = 1 or less</li> <li>MHCT V1 item 8 (Other) = 1 or less</li> </ul>	N/A



## CARE CLUSTER 2: Common mental health problems (low severity with greater need)

### Description:

This group has definite but minor problems of depressed mood, anxiety or other disorder but they do not present with any distressing psychotic symptoms. They may have already received care associated with cluster 1 and require more specific intervention, or previously been successfully treated at a higher level but are re-presenting with low level symptoms.

### Likely primary diagnosis:

Likely to include: F32 Depressive episode, F40 Phobic anxiety disorders, F41 Other anxiety disorders, F42 Obsessive-compulsive disorder, F43 Stress reaction/adjustment disorder, F50 Eating disorder.

### Unlikely primary diagnosis:

F00-03 Dementias, F20-29 Schizophrenia, schizotypal and delusional disorders, F30 Manic episode, F31 Bipolar disorder, F33 Major depressive disorder, recurrent.

### Impairment:

Disorder unlikely to cause disruption to wider functioning but some people will experience minor problems.

### Risk:

Unlikely to be an issue.

### Course:

The problem is likely to be short term and related to life events.

### Likely NICE guidance:

Service user experience in adult mental health CG136, Anxiety CG113, Depression in adults CG90, Depression with Chronic Health Problems CG91, Common mental health disorders CG123, OCD CG31, Eating Disorders CG9.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood*					
8	Other mental and behavioural problems*					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong Unreasonable Beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children and vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score

Unlikely to score

Expected to score

No data available

May score


\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 2: Common mental health problems

Indicative episode of care: 12 – 15 weeks

Cluster reviews at least every: 15 weeks

Cluster	Step-up criteria (The following criterion is met)	<u>Example local discharge Criteria</u> for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria (The following criterion is met)
2	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>MHCT V1 item 2 (Non-accidental self-injury) = 0</li> <li>MHCT V1 item 7 (Depression) = 1 or less</li> <li>MHCT V1 item 8 (Other) = 1 or less</li> </ul>	N/A



### CARE CLUSTER 3: Non-psychotic (moderate severity)

#### Description:

Moderate problems involving depressed mood, anxiety or other disorder (not including psychosis).

#### Likely primary diagnosis:

Likely to include: F32 Depressive episode, F40 Phobic anxiety disorders, F41 Other anxiety disorders, F42 Obsessive-compulsive disorder, F43 Stress reaction/adjustment disorder, F50 Eating disorder.

#### Unlikely primary diagnosis:

F00-03 Dementias, F20-29 Schizophrenia, schizotypal and delusional disorders, F30 Manic episode, F31 Bipolar disorder

#### Impairment:

Disorder unlikely to cause disruption to wider function but some people will experience moderate problems.

#### Risk:

Unlikely to be a serious issue.

#### Course:

Short-term.

#### Likely NICE guidance:

Service user experience in adult mental health CG136, Anxiety CG113, Depression in adults CG90, Depression with Chronic Health Problems CG91, Common mental health disorders CG123, OCD CG31, Eating Disorders CG9, Post-traumatic stress disorder (PTSD) CG 26.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children & vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score

Expected to score

May score

Unlikely to score

No data available


\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 3: Non-psychotic (moderate severity)

Indicative episode of care: 4-6 months

Cluster reviews at least every: 6 months

Cluster	Step-up criteria (The following criterion is met)	<u>Example local discharge criteria for MH services</u> ( <u>All</u> of the following criteria are met)	Step-down criteria ( <u>All</u> of the following criteria are met)
3	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>MHCT V1 item 2 (Non-accidental self-injury) = 0</li> <li>MHCT V1 item 7 (Depression) = 1 or less</li> <li>MHCT V1 item 8 (Other) = 1 or less</li> </ul>	<ul style="list-style-type: none"> <li>Patient has completed a successful period of treatment but is left with residual co-morbidities requiring an alternative treatment package at a lower intensity.</li> <li>Service user fits description and scoring profile of any likely/possible 'step-down' cluster</li> </ul>



## CARE CLUSTER 4: Non-psychotic (severe)

### Description:

The group is characterised by severe mood disturbance and/or anxiety and/or other increasing complexity of needs. They may experience disruption to function in everyday life and there is an increasing likelihood of significant risks.

### Likely primary diagnosis:

Likely to include: F32 Depressive episode, F40 Phobic anxiety disorders, F41 Other anxiety disorders, F42 Obsessive-compulsive disorder, F43 Stress reaction/adjustment disorder, F44 Dissociative disorder, F45 Somatoform disorder, F48 Other neurotic disorders, F50 Eating disorder.

### Unlikely primary diagnosis:

F00-03 Dementias, F20-29 Schizophrenia, schizotypal and delusional disorders, F30 Manic episode, F31.2&31.5 Bipolar disorder with psychosis

### Impairment:

Some may experience significant disruption in everyday functioning.

### Risk:

Some may experience moderate risk in self through self-harm or suicidal thoughts or behaviours.

### Course:

Unlikely to improve without treatment and may deteriorate with long term impact on functioning.

### Likely NICE guidance:

Service user experience in adult mental health CG136, Anxiety CG113, Depression in adults CG90, Depression with Chronic health Problems CG91, Common mental health disorders CG123, Medicines adherence CG76, OCD CG31, Eating Disorders CG9, Post-traumatic stress-disorder (PTSD) CG 26, Self-harm CG16.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children and vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score



Unlikely to score



Expected to score



No data available



May score




\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 4: Non-psychotic (severe)

Indicative episode of care: 6 - 12 months

Cluster reviews at least every: 6 months

Cluster	Step-up criteria (The following criterion is met)	<u>Example local discharge criteria for MH services</u> ( <u>All</u> of the following criteria are met)	Step-down criteria ( <u>All</u> of the following criteria are met)
4	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>MHCT V1 item 2 (Non-accidental self-injury) = 0</li> <li>MHCT V1 item 7 (Depression) = 1 or less</li> <li>MHCT V1 item 8 (Other) = 1 or less</li> </ul>	<ul style="list-style-type: none"> <li>Patient has completed a successful period of treatment but is left with residual co-morbidities requiring an alternative treatment package at a lower intensity.</li> <li>Service user fits description and scoring profile of any likely/possible 'step-down' cluster.</li> </ul>





## CARE CLUSTER 5: Non-psychotic disorders (very severe)

### Description:

This group will be experiencing severe mood disturbance and/or anxiety and/or other symptoms. They will not present with distressing hallucinations or delusions but may have some unreasonable beliefs. They may often be at high risk for non-accidental self-injury and they may present safeguarding issues and have severe disruption to everyday living.

### Likely primary diagnosis:

Likely to include: F32 Depressive episode (non-psychotic), F33 Recurrent depressive episode (non-psychotic), F40 Phobic anxiety disorders, F41 Other anxiety disorders, F42 Obsessive-compulsive disorder, F43 Stress reaction/adjustment disorder, F44 Dissociative disorder, F45 Somatoform disorder, F48 Other neurotic disorders, F50 Eating disorder.

### Unlikely primary diagnosis:

F00-03 Dementias, F20-29 Schizophrenia, schizotypal and delusional disorders, F30 Manic episode, F31.2 and 31.5 Bipolar disorder with psychosis

### Impairment:

Moderate or severe problems with relationships. Level of problems in other areas of role functioning likely to vary.

### Risk:

Likely moderate or severe risk of non-accidental self-injury with other possible risk, including safeguarding issues if any responsibility for young children or vulnerable dependent adults.

### Course:

Probably known to service for more than a year or expected to be known for an extended period.

### Likely NICE guidance:

Service user experience in adult mental health CG136, Anxiety CG113 Depression in adults CG90, Depression with Chronic health Problems CG91, Common mental health disorders CG123, Medicines adherence CG76, OCD CG31, Eating Disorders CG9, Post-traumatic stress-disorder (PTSD) CG 26, Self-harm CG16.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical Illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood*					
8	Other mental and behavioural problems*					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children and vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score



Unlikely to score



Expected to score



No data available



May score




\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 5: Non-psychotic (very severe)

Indicative episode of care: 1- 3 years.

Cluster reviews at least every: 6 Months

Cluster	Step-up criteria (The following criterion is met)	Example local discharge criteria for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria (The following criterion is met)
5	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>MHCT V1 item 2 (Non-accidental self-injury) = 0</li> <li>MHCT V1 item 7 (Depression) = 1 or less</li> <li>MHCT V1 item 8 (Other) = 1 or less</li> </ul>	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-down' cluster.</li> </ul>



## CARE CLUSTER 6: Non-psychotic disorder of over-valued ideas

Description:	No	ITEM DESCRIPTION	RATING				
			0	1	2	3	4
Moderate to very severe disorders that are difficult to treat. This may include mood disturbance treatment resistant eating disorder, OCD, etc. where extreme beliefs are strongly held, some personality disorders and enduring depression.	1	Overactive, aggressive, disruptive or agitated behaviour					
	2	Non-accidental self-injury					
<b>Likely primary diagnosis:</b> Likely to include: F32 Depressive episode (non-psychotic), F33 Recurrent Depressive Episode (non-psychotic), F40 Phobic Anxiety Disorders, F41 Other Anxiety Disorders, F42 Obsessive-compulsive disorder, F43 Stress reaction/adjustment disorder, F44 Dissociative disorder, F45 Somatoform disorder, F48 Other neurotic disorders, F50 Eating disorder and some F60.	3	Problem drinking or drug-taking					
	4	Cognitive problems					
<b>Unlikely primary diagnosis:</b> F00-03 Dementias, F20-29 Schizophrenia, schizotypal and delusional disorders, F30 Manic episode, F31.2&31.5 Bipolar disorder with psychosis	5	Physical illness or disability problems					
	6	Hallucinations and delusions					
<b>Impairment:</b> Likely to seriously affect activity and role functioning in many ways.	7	Depressed mood *					
	8	Other mental and behavioural problems *					
<b>Risk:</b> Unlikely to be a major feature but safeguarding may be an issue if any responsibility for young children or vulnerable dependant adults.	9	Relationships					
	10	Activities of daily living					
<b>Course:</b> The problems will be enduring.	11	Living conditions					
	12	Occupation and activities					
<b>Likely NICE guidance:</b> Service user experience in adult mental health CG136, Eating Disorders CG9, OCD CG31, Post-traumatic stress-disorder (PTSD) CG 26 Anxiety CG113, Depression in adults CG90, Medicines adherence CG76 Antisocial personality Disorder CG77, Borderline Personality Disorder CG78, Self-harm CG16, Self-harm (longer-term management) CG 133, Depression with Chronic health Problems CG91.	13	Strong unreasonable beliefs					
	A	Agitated behaviour/expansive mood					
	B	Repeat self-harm					
	C	Safeguarding other children & vulnerable dependant adults					
	D	Engagement					
	E	Vulnerability					
<b>Must score</b>							
<b>Expected to score</b>							
<b>May score</b>							
<b>Unlikely to score</b>							
<b>No data available</b>							


\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 6: Non-psychotic disorders of overvalued ideas

Indicative episode of care: 3 years +

Cluster reviews at least every: 6 months

Cluster	Step-up criteria (The following criterion is met)	<u>Example local discharge Criteria for MH services</u> ( <u>All</u> of the following criteria are met)	Step-down criteria (The following criterion is met)
6	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>MHCT V1 item 2 (Non-accidental self-injury) = 1 or less</li> <li>MHCT V1 item 7 (Depression) = 1 or less</li> <li>MHCT V1 item 8 (Other) = 1 or less</li> <li>MHCT V1 item 13 (Strong unreasonable beliefs) = 2 or less</li> </ul>	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-down' cluster.</li> </ul>



## CARE CLUSTER 7: Enduring non-psychotic disorders (high disability)

Description:	No	ITEM DESCRIPTION	RATING				
			0	1	2	3	4
This group suffers from moderate to severe disorders that are very disabling. They will have received treatment for a number of years and although they may improve in positive symptoms considerable disability remains that is likely to affect role functioning in many ways.	1	Overactive, aggressive, disruptive or agitated behaviour					
	2	Non-accidental self-injury					
	3	Problem drinking or drug-taking					
	4	Cognitive problems					
	5	Physical illness or disability problems					
	6	Hallucinations and delusions					
	7	Depressed mood *					
	8	Other mental and behavioural problems *					
	9	Relationships					
	10	Activities of daily living					
	11	Living conditions					
	12	Occupation and activities					
	13	Strong unreasonable beliefs					
Likely primary diagnosis: Likely to include: F32 Depressive episode (Non-Psychotic), F33 Recurrent depressive episode (non-psychotic), F40 Phobic anxiety disorders, F41 Other anxiety disorders, F42 Obsessive-compulsive disorder, F43 Stress reaction/adjustment disorder, F44 Dissociative disorder, F45 Somatoform disorder, F48 Other neurotic disorders, F50 Eating disorder and some F60.	A	Agitated behaviour/expansive mood					
	B	Repeat self-harm					
	C	Safeguarding other children and vulnerable dependant adults					
	D	Engagement					
	E	Vulnerability					
Unlikely primary diagnosis: F00-03 Dementias, F20-29 Schizophrenia, schizotypal and delusional disorders , F30 Manic episode, F31.2&31.5 Bipolar disorder with psychosis							
Impairment: Likely to seriously affect activity and role functioning in many ways.							
Risk: Unlikely to be a major feature but safeguarding may be an issue if any responsibility for young children or vulnerable dependant adults.							
Course: The problems will be enduring.							
Likely NICE guidance: Service user experience in adult mental health CG136, Eating Disorders CG9, OCD CG31, Anxiety CG113, Depression in adults CG90, Medicines adherence CG76, Post-traumatic stress-disorder (PTSD) CG26, Antisocial personality Disorder CG77, Borderline Personality Disorder CG78, Self-harm (longer-term management) CG 133, Depression with Chronic health Problems CG91							

Must score

Expected to score

May score

Unlikely to score

No data available


\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 7: Enduring non-psychotic disorders (high disability)

Indicative episode of care: 3 years +

Cluster reviews at least: Annually

Cluster	Step-up criteria (The following criterion is met)	Example local discharge criteria for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria (The following criterion is met)
7	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>MHCT V1 item 2 (Non-accidental self-injury) = 1 or less</li> <li>MHCT V1 item 7 (Depression) = 1 or less</li> <li>MHCT V1 item 8 (Other) = 1 or less N/A</li> </ul>	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/possible 'step-down' cluster.</li> </ul>



## CARE CLUSTER 8: Non-psychotic chaotic and challenging disorders

### Description:

This group will have a wide range of symptoms and chaotic and challenging lifestyles. They are characterised by moderate to very severe repeat deliberate self-harm and/or other impulsive behaviour and chaotic, over dependent engagement and often hostile with services.

### Likely primary diagnosis:

Likely to include F60 Personality disorder.

### Unlikely primary diagnosis:

F00-03 Dementias, F20-29 Schizophrenia, schizotypal and delusional disorders, F30 Manic episode, F31 Bipolar disorder.

### Impairment:

Poor role functioning with severe problems in relationships.

### Risk:

Moderate to very severe repeat deliberate self-harm, with chaotic, over dependent and often hostile engagement with service. Non-accidental self-injury risks likely to be present. Safeguarding may be an issue.

### Course:

The problems will be enduring.

### Likely NICE guidance:

Service user experience in adult mental health CG136, Borderline Personality Disorder CG78, Self-harm CG16, Self-harm (longer-term management) CG 133, Post-traumatic stress-disorder (PTSD) CG 26 Depression in adults CG90, Anxiety CG113, Alcohol dependence and harmful alcohol misuse CG115, Antisocial personality disorder CG77.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children & vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score



Unlikely to score



Expected to score



No data available



May score




\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 8: Non-psychotic chaotic and challenging disorders

Indicative episode of care: 3 years +

Cluster reviews at least: Annually

Cluster	Step-up criteria (The following criterion is met)	<u>Example local discharge criteria for MH services</u> ( <u>All</u> of the following criteria are met)	Step-down criteria (The following criterion is met)
8	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>MHCT V1 item 2 (Non-accidental self-injury) = 1 or less</li> <li>MHCT V1 item B (self-harm) = 1 or less</li> <li>MHCT V1 item 7 (Depression) = 1 or less</li> <li>MHCT V1 Item 8 (Other) = 1 or less</li> </ul>	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/possible 'step-down' cluster consistently for the past 12 months.</li> </ul>










## CARE CLUSTER 9: Blank cluster

<b>Description:</b>
<b>Likely primary diagnosis:</b>
<b>Unlikely primary diagnosis:</b>
<b>Impairment:</b>
<b>Risk:</b>
<b>Course:</b>
<b>Likely NICE guidance:</b>

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children and vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score		Unlikely to score	
Expected to score		No data available	
May score			

\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 9: Blank cluster

Indicative episode of care:

Cluster reviews at least:

Cluster	Step-up criteria (The following criterion is met)	<u>Example local discharge criteria</u> for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria (The following criterion is met)
	•	•	•

### Current Cluster 9

**Key:**

Most likely transition(s)

Possible transition

Rare Transition

## CARE CLUSTER 10: First episode psychosis (with/without manic features)

### Description:

This group will be presenting to the service for the first time with mild to severe psychotic phenomena. They may also have mood disturbance and/or anxiety or other behaviours. Drinking or drug-taking may be present but *will* not be the only problem.

### Likely primary diagnosis:

Likely to include (F20-F29) Schizophrenia, schizotypal and delusional disorders, F31 Bipolar disorder.

### Unlikely primary diagnosis:

F00-03 Dementias.

### Impairment:

Mild to moderate problems with activities of daily living. Poor role functioning with mild to moderate problems with relationships.

### Risk:

Vulnerable to harm from self or others. Some may be at risk of Non-accidental self-injury or a threat to others.

### Course:

First Episode.

### Likely NICE guidance:

Service user experience in adult mental health CG136, Schizophrenia (update) CG82, Bipolar disorder CG38, Medicines adherence CG76  
Depression in adults CG90, Anxiety CG113, Alcohol dependence and harmful alcohol misuse CG115, Self-Harm CG16.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children and vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score

Unlikely to score

Expected to score

No data available

May score


\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 10: First episode in psychosis

Indicative episode of care: 3 years.

Cluster reviews at least: Annually

Cluster	Step-up criteria (The following criterion is met)	Example local discharge criteria for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria ( <u>All</u> of the following criteria are met)
10	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>Requires no psychotropic medication or has been on a stable dose for the past year.</li> <li>Rates 0-1 on MHCT V1 item 6 (Hallucinations and Delusions)</li> <li>Not currently detained under the Mental Health Act.</li> <li>Has required no inpatient / IHT packages for the past year.</li> <li>Any residual risks can be managed by primary care.</li> <li>Rates 0-1 on MHCT V1 item 12 (Occupation and Activities).</li> <li>Level of social inclusion meets service user's expectations.</li> <li>Has received three years of intervention from an Early Intervention in Psychosis Team, or no longer feels they require a service.</li> </ul>	<ul style="list-style-type: none"> <li>Has a prescribed period of treatment from an Early Intervention in Psychosis Team or equivalent (depending on age).</li> <li>Service user fits description and scoring profile of any likely/ possible 'step down' cluster and the level of need is likely to be maintained until the next planned review.</li> </ul>



## CARE CLUSTER 11: Ongoing recurrent psychosis (low symptoms)

### Description:

This group has a history of psychotic symptoms that are currently controlled and causing minor problems if any at all. They are currently experiencing a sustained period of recovery where they are capable of full or near functioning. However, there may be impairment in self-esteem and efficacy and vulnerability to life.

### Likely primary diagnosis:

Likely to include (F20-F29) Schizophrenia, schizotypal and delusional disorders F30 Manic Episode, F31 Bipolar Affective Disorder.

### Unlikely primary diagnosis:

F00-03 Dementias, F32 Depressive episode, F33 Recurrent depressive disorder, F40-48 Neurotic, stress-related and somatoform disorders, F50 Eating disorders, F60 Specific personality disorders.

### Impairment:

Full or near full functioning.

### Risk:

Relapse.

### Course:

Long term.

### Likely NICE guidance:

Service user experience in adult mental health CG136, Schizophrenia (update) CG82, Bipolar disorder CG38.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical Illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children and vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score



Unlikely to score



Expected to score



No data available



May score




\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 11: Ongoing recurrent psychosis (low symptoms)

Indicative episode of care: 2 years +

Cluster reviews at least: Annually

Cluster	Step-up criteria (The following criterion is met)	Example local discharge criteria for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria (The following criterion is met)
11	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>Fits profile of cluster 11 at the point of the planned CPA review, and has done so consistently for the past 12 months.</li> <li>Requires no psychotropic medication or has been on a stable dose for the past year.</li> <li>Not currently detained under the Mental Health Act.</li> <li>Has required no inpatient / IHT packages for the past year.</li> <li>Any residual risks can be managed by primary care.</li> <li>Scores 0-1 on MHCT V1 item 12 (Occupation and Activities).</li> <li>Level of social inclusion meets service user's expectations.</li> </ul>	N/A



## CARE CLUSTER 12: Ongoing or recurrent psychosis (high disability)

### Description:

This group has a history of psychotic symptoms with a significant disability with major impact on role functioning. They are likely to be vulnerable to abuse or exploitation.

### Likely primary diagnosis:

Likely to include (F20-F29) Schizophrenia, schizotypal and delusional disorders F30 Manic episode, F31 Bipolar affective disorder.

### Unlikely primary diagnosis:

F00-03 Dementias, F32 Depressive episode, F33 Recurrent depressive disorder, F40-48 Neurotic, stress-related and somatoform disorders, F50 Eating disorders, F60 Specific personality disorders.

### Impairment:

Possible cognitive and physical problems linked with long-term illness and medication. May have limited survival skills and be lacking basic life skills and poor role functioning in all areas.

### Risk:

Vulnerable to abuse or exploitation.

### Course:

Long term.

### Likely NICE guidance:

Service user experience in adult mental health CG136, Schizophrenia (update) CG82, Bipolar disorder CG38, Self-Harm CG16, Self-harm (longer-term management) CG 133, Medicines adherence CG76.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children & vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score



Unlikely to score



Expected to score



No data available



May score




\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 12: Ongoing or recurrent psychosis (high disability)

Indicative episode of care: 3 years +

Cluster reviews at least: Annually

Cluster	Step-up criteria (The following criterion is met)	Example local discharge criteria for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria (The following criterion is met)
12	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>Fits profile of cluster 12 at the point of the planned CPA review, and has done so consistently for the past 12 months.</li> <li>Requires no psychotropic medication or has been on a stable dose for the past year.</li> <li>Not currently detained under the Mental Health Act.</li> <li>Has required no inpatient / IHT packages for the past year.</li> <li>Any residual risks can be managed by primary care.</li> <li>Scores 0-1 on MHCT V1 item 12 (Occupation and Activities).</li> <li>Level of social inclusion meets service user's expectations.</li> </ul>	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-down' cluster consistently for the past 12 months.</li> </ul>





## CARE CLUSTER 13: Ongoing or recurrent psychosis (high symptom and disability)

### Description:

This group will have a history of psychotic symptoms which are not controlled. They will present with severe to very severe psychotic symptoms and some anxiety or depression. They have a significant disability with major impact on role functioning.

### Likely primary diagnosis:

Likely to include (F20-F29) Schizophrenia, schizotypal and delusional disorders F30 Manic episode, F31 Bipolar affective disorder.

### Unlikely primary diagnosis:

F00-03 Dementias, F32 Depressive episode, F33 Recurrent depressive disorder, F40-48 Neurotic, stress-related and somatoform disorders, F50 Eating disorders, F60 Specific personality disorders

### Impairment:

Possible cognitive and physical problems linked with long-term illness and medication. May be lacking basic life skills and poor role functioning in all areas.

### Risk:

Vulnerability to abuse or exploitation.

### Course:

Long term.

### Likely NICE guidance:

Service user experience in adult mental health CG136, Schizophrenia (update) CG82, Bipolar disorder CG38, Medicines adherence CG76 Self-Harm CG16, Self-harm (longer-term management) CG 133.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children & vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score		Unlikely to score	
Expected to score		No data available	
May score			


\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 13: Ongoing or recurrent psychosis (high symptom and disability)

Indicative episode of care: 3 years +

Cluster reviews at least: Annually

Cluster	Step-up criteria (The following criterion is met)	<u>Example local discharge criteria</u> for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria (The following criterion is met)
13	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>Has received 2 years of specialist MH intervention.</li> <li>Requires no psychotropic medication or has been on a stable dose for the past year.</li> <li>Scores 0-1 on MHCT V1 item 6 (Hallucinations and Delusions)</li> <li>Not currently detained under the Mental Health Act.</li> <li>Has required no inpatient / IHT packages for the past year.</li> <li>Any residual risks can be managed by primary care.</li> <li>Scores 0-1 on MHCT V1 item 12 (Occupation and Activities).</li> <li>Level of social inclusion meets service user's expectations.</li> </ul>	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-down' cluster consistently for the past 12 months.</li> </ul>



## CARE CLUSTER 14: Psychotic crisis

### Description:

They will be experiencing an acute psychotic episode with severe symptoms that cause severe disruption to role functioning. They may present as vulnerable and a risk to others or themselves.

### Likely primary diagnosis:

Likely to include (F20-F29) Schizophrenia, schizotypal and delusional disorders F30 Manic episode, F31 Bipolar affective disorder.

### Unlikely primary diagnosis:

F00-03 Dementias, F32 Depressive episode, F33 Recurrent depressive disorder, F40-48 Neurotic, stress-related and somatoform disorders, F50 Eating disorders,

### Impairment:

Cognitive problems may present. Activities will be severely disrupted in most areas. Role functioning is severely disrupted in most areas.

### Risk:

There may be risks to self or others because of challenging behaviour and some vulnerability to abuse or exploitation. Also, possibly poor engagement with service. Safeguarding risk if parent/carer.

### Course:

Acute

### Likely NICE guidance:

Service user experience in adult mental health CG136, Schizophrenia (update) CG82, Bipolar disorder CG38, Medicines adherence CG76 Self-Harm CG16, Violence CG25.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children and vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score



Unlikely to score



Expected to score



No data available



May score




\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 14: Psychotic crisis

Indicative episode of care: 8 – 12 weeks

Cluster reviews at least every: 4 weeks

Cluster	Step-up criteria (The following criterion is met)	Example local discharge criteria for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria (The following criterion is met)
14	N/A	<ul style="list-style-type: none"> <li>Requires no psychotropic medication or has been on a stable dose and is adherent.</li> <li>Scores 0-1 on MHCT V1 item 6 (Hallucinations and Delusions).</li> <li>Any residual risks can be managed by Primary Care.</li> <li>Scores 0-2 on MHCT V1 item 12 (Occupation and Activities).</li> <li>Level of social inclusion meets service user's expectations.</li> </ul>	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-down' cluster.</li> </ul>



## CARE CLUSTER 15: Severe psychotic depression

### Description:

This group will be suffering from an acute episode of moderate to severe depressive symptoms. Hallucinations and delusions will be present. It is likely that this group will present a risk of non-accidental self-injury and have disruption in many areas of their lives.

### Likely primary diagnosis:

Likely to include, F32.3 Severe depressive episode with psychotic symptoms

### Unlikely primary diagnosis:

F00-03 Dementias, F40-48 Neurotic, stress-related and somatoform disorders, F50 Eating disorders,

### Impairment:

Cognitive problems may present. Activities will be severely disrupted in most areas. Role functioning is severely disrupted in most areas

### Risk:

Risk of Non-accidental self-injury and vulnerability likely to be present with other risks variable. Consider safeguarding risks if parent or carer.

### Course:

Acute

### Likely NICE guidance:

Service user experience in adult mental health CG136, Medicines adherence CG76, Depression in adults CG90, OCD CG31, Schizophrenia (update) CG82, Bipolar disorder CG38, Self-Harm CG16.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children and vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score		Unlikely to score	
Expected to score		No data available	
May score			


\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 15: Severe psychotic depression

Indicative episode of care: 8 – 12 weeks

Cluster reviews at least every: 4 weeks

Cluster	Step-up criteria (The following criterion is met)	<u>Example local discharge criteria</u> for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria (The following criterion is met)
15	N/A	<ul style="list-style-type: none"> <li>Requires no psychotropic medication or has been on a stable dose and is adherent.</li> <li>Scores 0-1 on MHCT V1 item 6 (Hallucinations and Delusions).</li> <li>Any residual risks can be managed by Primary Care.</li> <li>Scores 0-2 on MHCT V1 item 12 (Occupation and Activities).</li> <li>Level of social inclusion meets service user's expectations</li> </ul>	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-down' cluster.</li> </ul>



## CARE CLUSTER 16: Psychosis and affective disorder (high substance misuse and engagement)

### Description:

This group has enduring, moderate to severe psychotic or bipolar affective symptoms with unstable, chaotic lifestyles *and co-existing* problem drinking or drug-taking. They may present a risk to self and others and engage poorly with services. Role functioning is often globally impaired.

### Likely primary diagnosis:

Likely to include, (F10-F19) Mental and behavioural disorders due to psychoactive substance use (F20-F29) Schizophrenia, schizotypal and delusional disorders, Bipolar disorder

### Unlikely primary diagnosis:

F00-03 Dementias F32 Depressive episode, F33 Recurrent depressive disorder, F40-48 Neurotic, stress-related and somatoform disorders, F50 Eating disorders, F60 Specific personality disorders

### Impairment:

Physical Illness or disability problems may be present as a result of Problem drinking or drug-taking and possibly cognitively impaired as a consequence of psychotic features or problem drinking or drug-taking. Global impairment of role function likely.

### Risk:

Moderate to severe risk to other due to violent and aggressive behaviour. Likely to engage poorly with services. Some risk of accidental death.

### Course:

Long term.

### Likely NICE guidance:

Service user experience in adult mental health CG136, Psychosis with coexisting substance misuse CG120, Schizophrenia (update) CG82, Bipolar Disorder CG38, Medicines adherence CG76, Alcohol dependence and harmful alcohol misuse CG115, Alcohol Use Disorders CG100, Drug misuse-psychosocial interventions CG51, Drug-misuse – opioid detoxification CG100, Self-Harm CG16, Self-harm (longer-term management) CG 133.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children and vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score

Unlikely to score

Expected to score

No data available

May score


\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 16: Psychosis and affective disorder (high substance misuse and engagement)

Indicative episode of care: 3 years +

Cluster reviews at least every: 6 months

Cluster	Step-up criteria (The following criterion is met)	Example local discharge criteria for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria ( <u>All</u> of the following criteria are met)
16	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>Has received two years of specialist MH intervention.</li> <li>Requires no psychotropic medication or has been on a stable dose for the past year.</li> <li>Scores 0-1 on MHCT V1 item 6 (Hallucinations and delusions)</li> <li>Has required no inpatient / IHT packages for the past year.</li> <li>Any residual risks can be managed by primary care</li> <li>Scores 0-1 on MHCT V1 item 12 (Occupation and Activities).</li> <li>Level of social inclusion meets service user's expectations.</li> <li>Scores 0-1 MHCT V1 item D (Engagement).</li> </ul>	<ul style="list-style-type: none"> <li>Service user has fitted description and scoring profile of any likely/ possible 'step-down' cluster consistently for the past 12 months.</li> <li>Has required no inpatient / IHT packages for the past year.</li> <li>Scores 0-1 MHCT V1 item D (Engagement).</li> <li>Level of support (frequency of visits, etc) has been reduced to a level that can be provided by a less intensive care package for the past 6 months.</li> <li>MHCT V1 item 3 (Problem drinking or drug-taking) has remained at a score of 2 or less for the past 12 months</li> </ul>





## CARE CLUSTER 17: Psychosis and affective disorder – difficult to engage


Description: This group has moderate to severe psychotic symptoms with unstable, chaotic lifestyles. There may be some problems with drugs or alcohol not severe enough to warrant care associated with cluster 16. This group have a history of non-concordance, are vulnerable & engage poorly with services.	No	ITEM DESCRIPTION	RATING				
			0	1	2	3	4
Likely primary diagnosis: Likely to include, (F20-F29) Schizophrenia, schizotypal and delusional disorders, Bi-Polar	1	Overactive, aggressive, disruptive or agitated behaviour					
	2	Non-accidental self-injury					
	3	Problem drinking or drug-taking					
	4	Cognitive problems					
	5	Physical illness or disability problems					
Unlikely primary diagnosis: F00-03 Dementias, F32 Depressive episode, F33 Recurrent depressive disorder, F40-48 Neurotic, stress-related and somatoform disorders, F50 Eating disorders, F60 Specific personality disorders	6	Hallucinations and delusions					
	7	Depressed mood *					
	8	Other mental and behavioural problems *					
	9	Relationships					
Impairment: Possibly cognitively impaired as a consequence of psychotic features or Problem drinking or drug-taking including prescribed medication. Likely severe problems with relationships and one or more other area of functioning	10	Activities of daily living					
	11	Living conditions					
	12	Occupation and activities					
	13	Strong unreasonable beliefs					
Risk: Moderate to severe risk of harm to others due to aggressive or violent behaviour. Risk of non-accidental self-injury. Likely to be non-compliant, vulnerable and engage poorly with service.							
	A	Agitated behaviour/expansive mood					
	B	Repeat self-harm					
Course: Long term.	C	Safeguarding other children and vulnerable dependant adults					
	D	Engagement					
	E	Vulnerability					
Likely NICE guidance: Service user experience in adult mental health CG136, Schizophrenia (update) CG82, Bipolar Disorder CG38, Medicines adherence CG76 Alcohol Use Disorders CG100, Drug misuse-psychosocial interventions CG51, Psychosis with coexisting substance misuse CG120 Self-Harm CG16, Self-harm (longer-term management) CG 133.	<div> <div>Must score</div> <div>Unlikely to score</div> </div> <div> <div>Expected to score</div> <div>No data available</div> </div> <div> <div>May score</div> </div>						
	*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.						

## CARE TRANSITION PROTOCOLS - Cluster 17: Psychosis and affective disorder difficult to engage

Indicative episode of care: 3 years +

Cluster reviews at least every: 6 months

Cluster	Step-up criteria ( <u>Any</u> of the following criteria are met)	<u>Example local discharge criteria</u> for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria ( <u>All</u> of the following criteria are met)
17	<ul style="list-style-type: none"> <li>• Patient fits profile for clusters 14 or 15.</li> <li>• Patient scores above 2 on Problem drinking or drug-taking item and this results in an inability to deliver the care typically provided to cluster 17 patients without a significant increase in resources.</li> </ul>	<ul style="list-style-type: none"> <li>• Has received two years of specialist MH intervention.</li> <li>• Requires no psychotropic medication or has been on a stable dose for the past year.</li> <li>• Scores 0-1 on MHCT V1 item 6 (Hallucinations and delusions)</li> <li>• Has required no inpatient / IHT packages for the past year.</li> <li>• Any residual risks can be managed by primary care</li> <li>• Scores 0-1 on MHCT V1 item 12 (Occupation and Activities).</li> <li>• Level of social inclusion meets service user's expectations.</li> <li>• Scores 0-1 MHCT V1 item D (Engagement)</li> </ul>	<ul style="list-style-type: none"> <li>• Service user has fitted description and scoring profile of any likely/ possible 'step-down' cluster consistently for the past 12 months.</li> <li>• Has required no inpatient / IHT packages for the past year.</li> <li>• Scores 0-1 MHCT V1 item D (Engagement).</li> <li>• Level of support (frequency of visits, etc) has been reduced to a level that can be provided by a less intensive care package for the past six months.</li> </ul>



## CARE CLUSTER 18: Cognitive impairment (low need)

Description: People who may be in the early stages of dementia (or who may have an organic brain disorder affecting their cognitive function) who have some memory problems, or other low level cognitive impairment but who are still managing to cope reasonably well. Underlying reversible physical causes have been ruled out.	No	ITEM DESCRIPTION	RATING				
			0	1	2	3	4
Likely primary diagnosis: Diagnoses likely to include: F00 – Dementia in Alzheimer-s disease, F01 – Vascular dementia, F02 – Dementia in other diseases classified elsewhere F03 – Unspecified dementia, Dementia with Lewy bodies (DLB),	1	Overactive, aggressive, disruptive or agitated behaviour					
	2	Non-accidental self-injury					
	3	Problem drinking or drug-taking					
	4	Cognitive problems					
	5	Physical illness or disability problems					
	6	Hallucinations and delusions					
Unlikely primary diagnosis: F20-29 Schizophrenia, schizotypal and delusional disorders, F30-39 Mood [affective] disorders, F40-48 Neurotic, stress-related and somatoform disorders, F50 Eating disorders, F60 Specific personality disorders	7	Depressed mood *					
	8	Other mental and behavioural problems *					
	9	Relationships					
	10	Activities of daily living					
	11	Living conditions					
	12	Occupation and activities					
Impairment: Some memory and other low level impairment will be present. ADL function will be unimpaired, or only mildly impaired. There may be changes in ability to manage vocational and social roles.	13	Strong unreasonable beliefs					
Risk: None or minor.	A	Agitated behaviour/expansive mood					
	B	Repeat self-harm					
Course: Long term	C	Safeguarding other children and vulnerable dependant adults					
	D	Engagement					
Likely NICE guidance: Service user experience in adult mental health CG136, Dementia CG42 Medicines adherence CG76, Anxiety CG113, Depression in adults CG90, Depression with a chronic physical health problem CG91	E	Vulnerability					

Must score		Unlikely to score	
Expected to score		No data available	
May score			


\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 18: Cognitive impairment (low need)

Indicative episode of care: 3 years +

Cluster reviews at least every: Annually

Cluster	Step-up criteria (The following criterion is met)	Example local discharge criteria for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria ( <u>All</u> of the following criteria are met)
18	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-up' cluster.</li> </ul>	<ul style="list-style-type: none"> <li>Stable rating on MHCT item 4 (Cognitive problems) for the past year</li> <li>Stable dose of any prescribed ACEIs for the past six months</li> <li>Level of social inclusion meets service user's expectation</li> <li>Any residual risks can be managed by primary care</li> </ul>	<ul style="list-style-type: none"> <li>Organic causes of memory problems have been excluded.</li> <li>Service user fits description and scoring profile of any likely/ possible 'step-down' cluster.</li> </ul>



## CARE CLUSTER 19: Cognitive impairment or dementia complicated (moderate need)

Description: People who have problems with their memory, and or other aspects of cognitive functioning resulting in moderate problems looking after themselves and maintaining social relationships. Probable risk of self-neglect or harm to others and may be experiencing some anxiety or depression.	No	ITEM DESCRIPTION	RATING				
			0	1	2	3	4
Likely primary diagnosis: Likely to include: F00 – Dementia in Alzheimer's disease, F01 – Vascular dementia, F02 – Dementia in other diseases classified elsewhere, F03 – Unspecified dementia, F09 – unspecified organic or symptomatic mental disorder, Dementia with Lewy bodies (DLB), Frontotemporal dementia (FTD)	1	Overactive, aggressive, disruptive or agitated behaviour					
	2	Non-accidental self-injury					
	3	Problem drinking or drug-taking					
	4	Cognitive problems					
	5	Physical illness or disability problems					
	6	Hallucinations and delusions					
Unlikely primary diagnosis: F20-29 Schizophrenia, schizotypal and delusional disorders, F30-39 Mood [affective] disorders, F40-48 Neurotic, stress-related and somatoform disorders F50 Eating disorders, F60 Specific personality disorders	7	Depressed mood *					
	8	Other mental and behavioural problems *					
	9	Relationships					
	10	Activities of daily living					
	11	Living conditions					
	12	Occupation and activities					
Impairment: Impairment of ADL and some difficulty with communication and in fulfilling social and family roles.	13	Strong unreasonable beliefs					
Risk: Risk of self-neglect, harm to self or others. May lack awareness of problems.	A	Agitated behaviour/expansive mood					
	B	Repeat self-harm					
Course: Long term.	C	Safeguarding other children & vulnerable dependant adults					
	D	Engagement					
Likely NICE Guidance: Service user experience in adult mental health CG136, Dementia CG42 Medicines adherence CG76, Anxiety CG113, Depression in adults CG90 Depression with a chronic physical health problem CG91.	E	Vulnerability					

Must score

Expected to score

May score

Unlikely to score

No data available

\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 19: Cognitive impairment or dementia complicated (moderate need)

Indicative episode of care: 3 years +

Cluster reviews at least every: 6 months

Cluster	Step-up criteria (The following criterion is met)	<u>Example local discharge criteria</u> for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria ( <u>All</u> of the following criteria are met)
19	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible step up cluster</li> </ul>	<ul style="list-style-type: none"> <li>Stable rating on MHCT item 4 (Cognitive problems) for the past year</li> <li>Stable dose of any prescribed ACHEIs for the past 6 months</li> <li>No inpatient / home treatment packages for the last 12 months</li> <li>Level of social inclusion meets service user's expectation</li> <li>Any residual risks (including any comorbidities) can be managed by primary care</li> </ul>	N/A



## CARE CLUSTER 20: Cognitive impairment or dementia complicated (high need)

<b>Description:</b> People with dementia who are having significant problems in looking after themselves and whose behaviour may challenge their carers or services. They may have high levels of anxiety or depression, psychotic symptoms or significant problems such as aggression or agitation. They may not be aware of their problems. They are likely to be at high risk of self-neglect or harm to others, and there may be a significant risk of their care arrangements breaking down.	<b>No</b>	<b>ITEM DESCRIPTION</b>	<b>RATING</b>				
			<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
<b>Likely primary diagnosis:</b> Likely to include: F00 – Dementia in Alzheimer's disease, F01 – Vascular dementia, F02 – Dementia in other diseases classified elsewhere, F03 – Unspecified dementia, F09 – unspecified organic or symptomatic mental disorder, Dementia with Lewy bodies (DLB), Frontotemporal dementia (FTD).	1	Overactive, aggressive, disruptive or agitated behaviour					
	2	Non-accidental self-injury					
	3	Problem drinking or drug-taking					
	4	Cognitive problems					
	5	Physical illness or disability problems					
	6	Hallucinations and delusions					
	7	Depressed mood *					
	8	Other mental and behavioural problems *					
	9	Relationships					
	10	Activities of daily living					
	11	Living conditions					
	12	Occupation and activities					
	13	Strong unreasonable beliefs					
<b>Unlikely primary diagnosis:</b> F20-29 Schizophrenia, schizotypal and delusional disorders, F30-39 Mood [affective] disorders, F40-48 Neurotic, stress-related and somatoform disorders F50 Eating disorders, F60 Specific personality disorders							
	A	Agitated behaviour/expansive mood					
	B	Repeat self-harm					
	C	Safeguarding other children and vulnerable dependant adults					
	D	Engagement					
<b>Impairment:</b> Significant impairment of ADL function and/or communication. May lack awareness of problems. Significant impairment of role functioning. Unable to fulfil social and family roles	E	Vulnerability					
<b>Risk:</b> High risk of self-neglect or harm to self or others. Risk of breakdown of care.							
<b>Course:</b> Long term.							
<b>Likely NICE guidance:</b> Service user experience in adult mental health CG136, Dementia CG42 Medicines adherence CG76, Falls CG21, Anxiety CG113, Violence CG25 Depression in adults CG90, Depression with a chronic physical health problem CG91.							

<b>Must score</b>		<b>Unlikely to score</b>	
<b>Expected to score</b>		<b>No data available</b>	
<b>May score</b>			


**\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.**

## CARE TRANSITION PROTOCOLS - Cluster 20: Cognitive impairment or dementia complicated (high need)

Indicative episode of care: 3 years +

Cluster reviews at least every: 6 months

Cluster	Step-up criteria (The following criterion is met)	<u>Example local discharge criteria</u> for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria ( <u>All</u> of the following criteria are met)
20	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of and likely/ possible step up cluster.</li> </ul>	<ul style="list-style-type: none"> <li>Stable rating on MHCT item 4 (Cognitive problems) for the past 12 months</li> <li>Stable dose of any prescribed ACHEIs for the past 12 months</li> <li>No inpatient / home treatment packages for the last 12 months</li> <li>Level of social inclusion meets service users and carers expectation</li> <li>Any residual risks (including any comorbidities) can be managed by primary care with / without other partnerships</li> <li>Has received at least 1 year of specialist MH intervention</li> </ul>	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/possible 'step-down' cluster.</li> <li>Improvement is likely to be sustained until the next planned review</li> </ul>





## CARE CLUSTER 21: Cognitive impairment or dementia (high physical or engagement)

### Description:

People with cognitive impairment or dementia who are having significant problems in looking after themselves, and whose physical condition is becoming increasingly frail. They may not be aware of their problems and there may be a significant risk of their care arrangements breaking down.

### Likely primary diagnosis:

Likely to include: F00 – Dementia in Alzheimer's disease, F01 – Vascular dementia, F02 – Dementia in other diseases classified elsewhere, F03 – Unspecified dementia, F09 – unspecified organic or symptomatic mental disorder, Dementia with Lewy bodies (DLB), Frontotemporal dementia (FTD)

### Unlikely primary diagnosis:

F20-29 Schizophrenia, schizotypal and delusional disorders, F30-39 Mood [affective] disorders, F40-48 Neurotic, stress-related and somatoform disorders, F50 Eating disorders, F60 Specific personality disorders.

### Impairment:

Likely to lack awareness of problems. Significant impairment of ADL function. Unable to fulfil self-care and social and family roles. Major impairment of role functioning.

### Risk:

High risk of self-neglect. Risk of breakdown of care.

### Course:

Long term.

### Likely NICE guidance:

Service user experience in adult mental health CG136, Dementia CG42 Medicines adherence CG76, Falls CG21, Anxiety CG113.

No	ITEM DESCRIPTION	RATING				
		0	1	2	3	4
1	Overactive, aggressive, disruptive or agitated behaviour					
2	Non-accidental self-injury					
3	Problem drinking or drug-taking					
4	Cognitive problems					
5	Physical illness or disability problems					
6	Hallucinations and delusions					
7	Depressed mood *					
8	Other mental and behavioural problems *					
9	Relationships					
10	Activities of daily living					
11	Living conditions					
12	Occupation and activities					
13	Strong unreasonable beliefs					
A	Agitated behaviour/expansive mood					
B	Repeat self-harm					
C	Safeguarding other children and vulnerable dependant adults					
D	Engagement					
E	Vulnerability					

Must score



Unlikely to score



Expected to score



No data available



May score




\*Use the highest rating from Scales 7 and 8 when deciding if the rating fits the range indicated.

## CARE TRANSITION PROTOCOLS - Cluster 21: Cognitive impairment or dementia (high physical or engagement needs)

Indicative episode of care: 3 years +

Cluster reviews at least every: 6 months

Cluster	Step-up criteria (The following criterion is met)	<u>Example local discharge criteria</u> for MH services ( <u>All</u> of the following criteria are met)	Step-down criteria ( <u>All</u> of the following criteria are met)
21	N/A	<ul style="list-style-type: none"> <li>No inpatient / home treatment packages for the last 12 months</li> <li>Level of social inclusion meets Service Users and carers expectation</li> <li>Any residual risks (including any comorbidities with the use of the Principles of Palliative Care Approach / Specialist Palliative care) can be managed by primary care with / without other partnerships</li> <li>Has received at least 1 year of specialist MH intervention</li> </ul>	<ul style="list-style-type: none"> <li>Service user fits description and scoring profile of any likely/ possible 'step-down' cluster.</li> <li>Improvement is likely to be sustained until the next planned review</li> </ul>



## Appendix 4 – Practice guidance note regarding: allocation of patients with a bipolar diagnosis to a cluster

### Summary

Following requests for extra guidance about the allocation of patients with a diagnosis of bipolar disorder to the nationally mandated needs-based clusters, the following statement has been produced.

**All patients, including those with a diagnosis of bipolar disorder, should be allocated to the cluster that **best describes the combination and severity of their primary presenting needs**. Patients with the same diagnosis can therefore be accurately allocated to different clusters within a superclass (non-psychotic, psychotic/organic). As a diagnosis of bipolar disorder covers a particularly wide variety of presentations these patients may be allocated to either a psychotic or a non-psychotic cluster depending on your clinical judgement, though consideration of the likely and unlikely diagnoses sections of the clustering booklet should be carefully considered.**

See below for a fuller rationale.

### Background

The 21 mandated clusters were originally designed to be a purely needs-led classification system. The concept of diagnosis was included retrospectively as diagnostic labels were a helpful way to succinctly describe superclasses. Likely primary diagnoses were also included to assist medical staff to understand how the model aligned to their diagnostic classification system.

More recently 'unlikely primary diagnoses' were included, primarily to help with retrospective audits of clustering accuracy. It should be noted that the low levels of diagnosis recording nationally meant that the lists of unlikely diagnoses by cluster were initially generated from a single provider organisation's data and then moderated by wider clinical opinion.

The cumulative effect of these developments has incrementally led to some stakeholders wanting to create a 1:1 relationship between cluster and diagnosis. This has become a particular issue for the diagnosis of bipolar disorder where there is perhaps the most tension between a diagnostic approach and a needs-led classification system.

## Bipolar diagnosis

It is well recognised that a single diagnosis can span a number of clusters, depending on severity. In the case of bipolar affective disorder, the range of possible presentations makes this particularly likely (see below) as does the distinction between bipolar I and bipolar II (though the latter is yet to appear in ICD 10).

## Current state

The clustering process has now been used nationally for some time and queries continue to be raised regarding patients diagnosed with bipolar disorder, and in particular, those with the Diagnostic and Statistical Manual of Mental Disorders (IV) (DSM-iv) definition of bipolar II.

Differing local guidance has been established and so, in addition to the wide range of presentations described above, there is an equally wide range of stakeholders and opinions. These range from those who continue to seek a 1:1 relationship between cluster and diagnosis to those who would seek to completely dissociate the two approaches.

Our advice is that those with a diagnosis of bipolar disorder should be allocated to the cluster that **best describes the combination and severity of their primary presenting needs**. Patients with the same diagnosis can therefore be accurately allocated to different clusters within a superclass (non-psychotic, psychotic / organic). However, someone with bipolar disorder should not normally be assigned to clusters 1-4, as the indicative episode of care is likely to be at odds with the course of bipolar disorder.

## Current generic guidance

The current national guidance for allocation to cluster is:

- **Step 1:** Based on the information you have gathered during your routine screening/assessment process, rate the individual's identified needs using the Mental Health Clustering Tool – Version 3.0 (Appendix 1).
- **Step 2:** Use the Decision Tree (Appendix 2) to decide if the presenting needs are non-psychotic, psychotic or organic in origin. Then decide which of the next level of headings is most accurate. This will narrow down the list of clusters likely to describe the person's needs.
- **Step 3:** Look at the rating grids (Appendix 3) to decide which one is the most appropriate by using the colour-coded key.

- Start with the red ratings. These indicate the type and level of need that must be rated. If the ratings do not match, try another cluster.
- Next, consider the orange ratings. These represent expected ratings. You may allocate a person to a cluster if the orange ratings do not exactly match the coloured grids. However, this reflects a ‘weaker fit’ to that cluster.
- Finally review the yellow ratings. These represent ratings that may occur. These scales have significantly less bearing on cluster allocation but may indicate the need for extra care plan interventions.

**Remember, the final clustering decision is yours, based on your assessment results and your clinical judgement in applying this guidance.**

### Case vignettes

The following vignettes are intended to illustrate the application of this guidance:

#### *Bipolar 1 vignette – psychosis pathway: ‘Harry’*

Harry is a 39-year-old, single man. He is the eldest of 9 siblings and was fostered from the age of 5 until he was 11, when he was returned to his family. Harry’s natural mother was very strict and beat him with a leather belt for minor infractions of her rules. He is close to some of his siblings but no longer has any contact with either biological parent.

Harry had conduct problems at school. He left without any qualifications aged 15 and shortly afterwards he left home. Harry first presented mental health services at the age of 19 and for many years he was admitted to hospital formally on average 2-3 times each year. For the past 10 years the frequency of his admissions has decreased but he has a formal admission at least annually and he often requires psychiatric intensive care unit (PICU) management. Harry has received numerous, intensive psychological treatment packages without lasting impact. He is now managed on a community treatment order which enables his clinical team to intervene at an early stage when he relapses.

Harry lives alone in housing association accommodation and a housing support worker helps him to manage the tenancy. He has a poor history of retaining tenancies due to antisocial behaviour when unwell. Harry has a history of drug and alcohol misuse and a history of assault on family members, neighbours, girlfriends, other patients and members of staff. He has attempted suicide on several occasions.

When Harry is well he presents as a charming, engaging and gregarious man. His personal care is good and he often wears designer clothes. He is fully independent with activities of daily living and takes care of his tenancies but often gets into difficulty with neighbours due to noise nuisance. In the past he has been persuaded to spend his time constructively at day centres and educational/training centres.

When Harry relapses he poses a risk to himself and others. His relapse signature includes defaulting on his mood stabiliser and other medication, often shortly after leaving hospital, and then his mood drops. He then uses street drugs as 'pick me ups' and his mental state deteriorates rapidly.

When he is unwell Harry generally presents in a manic state with pressured speech and flights of ideas. He is elated, grandiose and sexually disinhibited. Harry is thought disordered, hostile, agitated and aggressive when unwell. He experiences auditory hallucinations and has paranoid ideas. He denies thought insertion, withdrawal, broadcast or passivity phenomena. His diagnosis has been reviewed on many occasions but bipolar disorder is the most consistent and enduring diagnosis over the years. When Harry relapses he is very vulnerable and has been observed dancing in the traffic on a busy road and then sustaining cuts and bruises when a car collided with him.

Harry is managed on a psychosis care pathway and he is assigned to Care Cluster 16 reflecting his complex presentation, co-morbidities and MHCT scores at assessment. When Harry relapses he is assigned to Cluster 14 for the duration of the acute psychotic crisis before stepping back down to Cluster 16.

#### *Bipolar 2 vignette – non-psychosis pathway: 'Elaine'*

Elaine is a 29-year-old single woman who lives at home with her parents. She works as an accountant in the City of London. Her parents describe Elaine's pre-morbid personality as 'shy' and 'lacking self-confidence'. Elaine does not socialise very much but enjoys the company of a small network of friends and work colleagues. She has never misused drugs or alcohol.

Elaine first came into contact with secondary mental health services at the age of 23 when she was preparing to sit an accountancy exam. Her parents observed that Elaine was unusually cheerful for no apparent reason and was waking very early. She disturbed her parents by making coffee at five in the morning while singing along to the radio. When her parents complained that Elaine was disturbing their sleep she became irritable and verbally aggressive, which was quite out of character.

During this period, lasting approximately two weeks, Elaine surprised her parents by returning home wearing garish, expensive clothes she had bought in the West End.

Elaine's work colleagues also noted the change in her demeanour. She was unusually sociable, chatty and enthusiastic about everything. Elaine told them she was destined to succeed in life and was expecting a senior promotion and a substantial salary increase. Her colleagues played along, enthralled by her optimism and unusually witty conversation. Elaine's boss noticed she would spontaneously initiate intense conversations with staff she scarcely knew in adjoining offices and he also noticed a decline in the quality of her work.

One evening after work Elaine attended a leaving party for a colleague at a nearby hotel. As the evening developed Elaine's speech became louder, more pressured and incoherent. She embarrassed several elderly male colleagues by trying to seduce them in a loud, provocative and sexually disinhibited manner. The hotel receptionist called the police after Elaine proposed they have sex, and ignoring his polite refusal tried repeatedly to grope his genitals. Elaine came into hospital informally with the support of her parents. There followed a rapid, dramatic reversal of mood and she became severely depressed and suicidal. Elaine experienced a similar hypomanic episode two years later at the age of 25 in the context of a relationship breakdown.

For the past four years there have been no further hypomanic episodes. Elaine continues to work as an accountant but with a different city firm which has positive policies regarding the employment of people with mental health problems. Elaine receives ongoing psychological and medical support from her local community mental health team. Elaine sporadically experiences brief episodes when she feels intensely happy, highly energised and optimistic about the future but these episodes are interspersed with much longer periods of mild to moderate depression. She has no psychotic experiences and she complains that 'I never get high any more'.

Elaine is now managed on a non-psychosis care pathway. She was assigned to Cluster 5 when severely depressed. Her Care Co-ordinator has now stepped down to a Cluster 7 care package, reflecting Elaine's MHCT scores, the improvement in positive symptoms and the duration/chronicity of her presentation.

Contact us:

**NHS Improvement**

Wellington House

133-155 Waterloo Road

London

SE1 8UG

**0300 123 2257**

**[pricing@improvement.nhs.uk](mailto:pricing@improvement.nhs.uk)**

**improvement.nhs.uk**

NHS Improvement publication code: CG 34/19

NHS England Publishing Approval Reference: 000298